arXiv:1411.6398v1 [math.OA] 24 Nov 2014

Nonlinear completely positive maps and dilation theory for real involutive algebras

Daniel Beltiţă and Karl-Hermann Neeb

24.11.2014

Abstract

A real seminormed involutive algebra is a real associative algebra $\mathcal{A}$ endowed with an involutive antiautomorphism $*$ and a submultiplicative seminorm p with $p(a^*) = p(a)$ for $a \in \mathcal{A}$. Then $\mathtt{ball}(\mathcal{A}, p) := \{a \in \mathcal{A} \colon p(a) < 1\}$ is an involutive subsemigroup. For the case where $\mathcal{A}$ is unital, our main result asserts that a function $\varphi \colon \mathtt{ball}(\mathcal{A}, p) \to B(V)$, V a Hilbert space, is completely positive (defined suitably) if and only if it is positive definite and analytic for any locally convex topology for which $\mathtt{ball}(\mathcal{A}, p)$ is open. If $\eta_{\mathcal{A}} \colon \mathcal{A} \to C^*(\mathcal{A}, p)$ is the enveloping C^*-algebra of $(\mathcal{A}, p)$ and $e^{C^*(\mathcal{A},p)}$ is the c_0-direct sum of the symmetric tensor powers $S^n(C^*(\mathcal{A}, p))$, then the above two properties are equivalent to the existence of a factorization $\varphi = \Phi \circ \Gamma$, where $\Phi \colon e^{C^*(\mathcal{A},p)} \to B(V)$ is linear completely positive and $\Gamma(a) = \sum_{n=0}^{\infty} \eta_{\mathcal{A}}(a)^{\otimes n}$. We also obtain a suitable generalization to non-unital algebras.

An important consequence of this result is a description of the unitary representations of $\mathrm{U}(\mathcal{A})$ with bounded analytic extensions to $\mathtt{ball}(\mathcal{A}, p)$ in terms of representations of the C^*-algebra $e^{C^*(\mathcal{A},p)}$.

Mathematics Subject Classification 2000: 22E65, 46L05, 46L07

Keywords and phrases: C^*-algebra, $*$-semigroup, involutive algebra, completely positive map, unitary group

Contents

1 Introduction **2**

2 Basic concepts **5**

3 Positive definite functions and dilations **10**
3.1 Preliminaries on positive definite functions 10
3.2 Multipliers and structured $*$-semigroups 14
3.3 Extreme points and homogeneity 20

4 Completely positive functions on $\mathtt{ball}(\mathcal{A}, p)$ **22**
4.1 Complete positivity of dilations 22
4.2 The case $\mathcal{A} = \mathbb{R}$ 26
4.3 Series expansion of bounded completely positive functions 27

5 Analytic positive definite functions **28**
5.1 Series expansions on structured $*$-semigroups 29
5.2 Analytic positive definite functions 30
5.3 Homogeneous positive definite functions 32

6 Proof of the main theorem **36**

7 Relations to Arveson's c.p. concept **39**

A Analytic functions on infinite dimensional spaces **41**

1 Introduction

A real *seminormed involutive algebra (or $*$-algebra)* is a pair $(\mathcal{A}, p)$, consisting of a real associative algebra $\mathcal{A}$ endowed with an involutive antiautomorphism $*$ and a submultiplicative seminorm p satisfying $p(a^*) = p(a)$ for $a \in \mathcal{A}$. Our project eventually aims at a systematic understanding of unitary representations of unitary groups $\mathrm{U}(\mathcal{A})$ of real seminormed involutive algebras $(\mathcal{A}, p)$. If $\mathcal{A}$ is unital, its unitary group is

$$\mathrm{U}(\mathcal{A}) := \{a \in \mathcal{A} : a^*a = aa^* = \mathbf{1}\},$$

and if $\mathcal{A}$ is non-unital, then $\mathrm{U}(\mathcal{A}) := \mathrm{U}(\mathcal{A}^1) \cap (\mathbf{1} + \mathcal{A})$, where $\mathcal{A}^1 = \mathcal{A} \oplus \mathbb{R}\mathbf{1}$ is the unitization of $\mathcal{A}$. Typical examples we have in mind are C^*-algebras (considered as real algebras) and algebras of the form $\mathcal{A} = C^\infty(X, \mathcal{B})$ for a Banach $*$-algebra $\mathcal{B}$ or $\mathcal{A} = C^\infty(X, M_n(\mathbb{K}))$, where X is a smooth manifold and $\mathbb{K} \in \{\mathbb{R}, \mathbb{C}, \mathbb{H}\}$. In the latter case $\mathrm{U}(\mathcal{A}) \cong C^\infty(X, \mathrm{U}_n(\mathbb{K}))$ are groups of smooth maps with values in a compact Lie group, hence particular examples of gauge groups.

We wish to initiate a line of investigation of unitary representations of $\mathrm{U}(\mathcal{A})$ which, in a suitable sense, are boundary values of representations of the ball semigroups $\mathtt{ball}(\mathcal{A}) := \{a \in \mathcal{A} \colon p(a) < 1\}$. This leads us naturally to completely positive maps on these semigroups. The present paper provides the foundation of that study, by developing the corresponding dilation theory and reducing it completely to the C^*-context. In a sequel we will apply these dilation methods to the representation theory of semigroups $\mathtt{ball}(\mathcal{A}, p)$ for various types of concrete algebras $\mathcal{A}$.

We now introduce some concepts needed to state our main theorem. If S is an involutive semigroup and V a Hilbert space, then a positive definite function $\varphi \colon S \to B(V)$ is called *dilatable* if there exist a representation $(\pi, \mathcal{H})$ of S and a continuous linear map $\iota \colon V \to \mathcal{H}$ such that $\varphi(s) = \iota^*\pi(s)\iota$ for $s \in S$. We write $\eta_{\mathcal{A}} \colon \mathcal{A} \to C^*(\mathcal{A}, p)$ for the enveloping C^*-algebra of the real seminormed $*$-algebra $(\mathcal{A}, p)$, and for a C^*-algebra $\mathcal{B}$, we write $e^{\mathcal{B}}$ for the c_0-direct sum of the C^*-algebras $S^n(\mathcal{B}) \subseteq \mathcal{B}^{\otimes n}$, where the tensor products are constructed from the maximal C^*-cross norm (see [Arv87] and Definition 6.1 below).

Here is our main theorem:

Theorem 1.1. *Let $(\mathcal{A}, p)$ be a real seminormed involutive algebra and V a complex Hilbert space. For a bounded function $\varphi \colon \mathtt{ball}(\mathcal{A}, p) \to B(V)$, the following are equivalent:*

(i) *φ is completely positive and dilatable.*

(ii) *φ is dilatable, positive definite and analytic with respect to some locally convex topology for which p is continuous.*

(iii) *There exists a linear completely positive map $\Phi\colon e^{C^*(\mathcal{A},p)} \to B(V)$ with $\Phi \circ \Gamma = \varphi$, where $\Gamma(a) = \sum_{n=0}^{\infty} \eta_{\mathcal{A}}(a)^{\otimes n}$.*

If $\mathcal{A}$ is unital, then every bounded positive definite function on $\mathtt{ball}(\mathcal{A}, p)$ is dilatable, so that this requirement can be omitted in (i) *and* (ii).

Note that (ii) is particularly strong because analyticity is not required for the topology defined by p but for some topology which may be much finer. An important consequence of this theorem is that it leads to a one-to-one correspondence between bounded analytic representations of $\mathtt{ball}(\mathcal{A}, p)$ with the representations of the C^*-algebra $e^{C^*(\mathcal{A},p)}$ (Theorem 6.8), which thus plays the role of a host algebra in the sense of [Gr05] for the bounded analytic representations of the involutive semigroup $\mathtt{ball}(\mathcal{A}, p)$, resp., the corresponding unitary representations of $\mathrm{U}(\mathcal{A})$, obtained from its action on $\mathtt{ball}(\mathcal{A}, p)$ by multipliers.

To see how such representations arise naturally, consider the unitary group $\mathrm{U}(\mathcal{A})$ of a C^*-algebra $\mathcal{A}$. Then every irreducible representation $(\pi, \mathcal{H})$ of $\mathcal{A}$ provides for every partition $\lambda = (\lambda_1, \dots, \lambda_k)$ of $N \in \mathbb{N}$ an irreducible representation $(\pi_\lambda, \mathbb{S}_\lambda(\mathcal{H}))$ on a subspace of $\mathcal{H}^{\otimes N}$ by a straightforward generalization of the classical Schur–Weyl theory to Hilbert spaces (see [BN12], and [EI13, Nes13] for an extension to type II_1-factor representations). Any such representation extends to a (holomorphic) polynomial representation of the multiplicative semigroup $(\mathcal{A}, \cdot)$, resp., the C^*-algebra $S^N(\mathcal{A})$ on $\mathbb{S}_\lambda(\mathcal{H})$.

In the special case where $\mathcal{A} = K(\mathcal{H})$ is the C^*-algebra of compact operators on an infinite dimensional complex Hilbert space $\mathcal{H}$, results of A. Kirillov and G. Olshanski [Ki73, Ol78] provide a complete classification of all unitary representation of $\mathrm{U}(\mathcal{A})$. All continuous unitary representations of $\mathrm{U}(\mathcal{A})$ are direct sums of irreducible ones which are of the form $\mathbb{S}_\lambda(\mathcal{H}) \otimes \mathbb{S}_\mu(\mathcal{H})^*$ and there are natural generalizations to $\mathrm{U}(K(\mathcal{H}))$, where $\mathcal{H}$ is a real or a quaternionic Hilbert space. Therefore the Schur–Weyl construction is exhaustive in these three cases. The key method to obtain these results is to show that continuous unitary representations of $\mathrm{U}(K(\mathcal{H}))$ are generated by dilation from operator-valued positive definite functions φ of the form $\varphi(g) = \rho(ege)$, where e is a hermitian projection of finite rank and ρ is a continuous representation of the semigroup $S = \overline{\mathtt{ball}}(e\mathcal{A}e)$ of contractions in $e\mathcal{A}e$, where $e\mathcal{A}e \cong M(n, \mathbb{K})$ for $\mathbb{K} \in \{\mathbb{R}, \mathbb{C}, \mathbb{H}\}$. Comparing with the results of the present paper, it follows that the representations ρ of S for which φ is positive definite are precisely the completely positive ones (see [Ne13] for a recent presentation of the Olshanski–Kirillov theory from this point of view). We conclude that, in this case, $e^{C^*(K(\mathcal{H}))} \cong e^{K(\mathcal{H}_{\mathbb{C}})}$ is a host algebra for all continuous unitary representations of $\mathrm{U}(K(\mathcal{H}))$.

For the algebra $\mathcal{A} = C^\infty(X, M_n(\mathbb{K}))$, X a compact smooth manifold, the identity component of $\mathrm{U}(\mathcal{A})$ is a product of an abelian group and the group $C^\infty(X, \mathrm{SU}_N(\mathbb{K}))_0$ whose norm continuous unitary representations have recently been classified ([JN13]). Here the irreducible ones are finite tensor products of so-called evaluation representations, hence obtained by restricting finite tensor products of irreducible algebra representations. In [BN15], we will see how all these particular results fit naturally into our present framework and how the representations of $S^n(\mathcal{B})$ for a C^*-algebra $\mathcal{B}$ can be described in terms of representations of $\mathcal{B}$.

For the proof of our main theorem, we need several results that are interesting in their own right:

- Factorization of $*$-representations of degree n of $(\mathcal{A}, \cdot)$ bounded on $\mathtt{ball}(\mathcal{A}, p)$ through a linear representations of the C^*-algebra $S^n(C^*(\mathcal{A}, p))$. So the C^*-functor is compatible with this kind of non-linearity.

- If φ is completely positive and homogeneous of degree $\alpha \in \mathbb{R}$, then $\alpha \in \mathbb{N}_0$ and φ is polynomial. This is based on Arveson's method of iterated differences ([Arv87]) and the case $\mathcal{A} = \mathbb{R}$, for which the completely positive maps $\varphi\colon \overline{\mathtt{ball}}(\mathcal{A}) = [-1, 1] \to \mathbb{R}$ are determined in [CR78].

Theorem 1.1 generalizes Arveson's work on non-linear states on balls of unital C^*-algebras ([Arv87]) which we found very inspiring. Actually he uses a notion of complete positivity that is weaker than ours and does not imply positive definiteness although he uses it at some point in his arguments. We discuss this issue in Section 7. His result asserts that completely positive states $\varphi\colon \mathtt{ball}(\mathcal{A}, p) \to \mathbb{C}$ factor through linear states of the C^*-algebra $e^{\mathcal{A}} \otimes e^{\overline{\mathcal{A}}}$. This mysterious occurrence of the C^*-algebra $\overline{\mathcal{A}}$ is clarified by our present approach via real algebras in which $C^*(\mathcal{A}) \cong \mathcal{A}_{\mathbb{C}} \cong \mathcal{A} \oplus \overline{\mathcal{A}}$, so that $e^{C^*(\mathcal{A})} \cong e^{\mathcal{A}} \otimes e^{\overline{\mathcal{A}}}$ provides the bridge to our Theorem 1.1(iii).

Theorem 1.1 also extends work of Ando/Choi on expanding nonlinear completely positive maps $\varphi\colon \mathcal{A} \to \mathcal{B}$ between C^*-algebras as series $\varphi = \sum_{mn,=0}^{\infty} \varphi_{m,n}$, where the maps $\varphi_{m,n}$ are homogeneous polynomial and completely positive. For the special case case where $\mathcal{A}$ is a C^*-algebra, the equivalence of (i) and (iii) in Theorem 1.1 is due to Hiai/Nakamura ([HN87]). If $\mathcal{A}$ is a C^*-algebra, then the situation simplifies considerably because an application of Arveson's Extension Theorem ([Arv69, Thm. 1.2.3]) shows that every bounded completely positive map φ on $\mathtt{ball}(\mathcal{A}, p)$ is dilatable.

The structure of this paper is as follows. In Section 2 we introduce the basic concepts, such as real seminormed involutive algebras, their ball semigroups and completely positive functions thereon. Section 3 develops dilation theory on the abstract level of structured $*$-semigroups. These are involutive semigroups S with a homomorphism $\gamma\colon (0, 1] \to S$, so that $S^\circ := \bigcup_{0<r<1} \gamma(r)S$ is a semigroup ideal. Here our main result is Theorem 3.18 on the existence of dilations for positive definite functions on S°. This applies in particular to the subsemigroup $S^\circ = \mathtt{ball}(\mathcal{A}, p)$ for the structured $*$-semigroup $S = \overline{\mathtt{ball}}(\mathcal{A}, p)$, when $\mathcal{A}$ is unital. In Section 4 we provide the key tools to obtain a series expansion $\varphi = \sum\limits_{n=0}^{\infty} \varphi_n$ of a bounded completely positive function φ on $\mathtt{ball}(\mathcal{A}, p)$ into homogeneous components φ_n.

In Section 5 we turn to positive definite functions $\varphi\colon \mathtt{ball}(\mathcal{A}, p) \to B(V)$ which are analytic with respect to some locally convex topology on $\mathcal{A}$ for which $\mathtt{ball}(\mathcal{A}, p)$ is open. We derive a series expansion of dilatable bounded analytic positive definite functions, but to use this result to show that φ is completely positive, a detailed analysis of the case where φ is a homogeneous polynomial is required. Here the isomorphism $C^*(S^n(\mathcal{A}), p_n) \cong S^n(C^*(\mathcal{A}, p))$ is a key ingredient. All these partial results are combined in Section 6 to a proof of Theorem 1.1.

2 Basic concepts

In this section we introduce real seminormed involutive algebras $(\mathcal{A}, p)$ and their enveloping C^*-algebras $C^*(\mathcal{A}, p)$. We further introduce completely positive maps on these algebras and their open unit balls. We conclude this section with a discussion of complexifications and embeddings of non-unital algebras into unital ones and how this is reflected by the enveloping C^*-algebra.

Seminormed involutive algebras

Definition 2.1. A *real involutive algebra* is a real associative algebra $\mathcal{A}$, endowed with an involutive antiautomorphism $a \mapsto a^*$.

(a) A submultiplicative seminorm $p\colon \mathcal{A} \to [0,\infty)$ is called *involutive* if $p(a^*) = p(a)$ for all $a \in \mathcal{A}$. If $\mathcal{A}$ is unital, we further assume $p(\mathbf{1}) = 1$. A *seminormed involutive algebra* is a pair $(\mathcal{A}, p)$, consisting of an involutive algebra $\mathcal{A}$ and an involutive submultiplicative seminorm p on $\mathcal{A}$ (cf. [Ne00, Def. III.2.3]).

(b) We will use the following notation:

- $\mathrm{U}(\mathcal{A}) := \{u \in \mathcal{A} \mid u^*u = uu^* = \mathbf{1}\}$ for the *unitary group* of a unital $*$-algebra $\mathcal{A}$ and for a non-unital algebra we put $\mathrm{U}(\mathcal{A}) := \mathrm{U}(\mathcal{A}^1) \cap (\mathbf{1} + \mathcal{A})$, where $\mathcal{A}^1 = \mathcal{A} \oplus \mathbb{R}\mathbf{1}$ is the unitization of $\mathcal{A}$;
- $\mathtt{ball}(\mathcal{A}, p) := \{a \in \mathcal{A} \mid p(a) < 1\}$ for the *open ball semigroup* of $(\mathcal{A}, p)$;
- $\overline{\mathtt{ball}}(\mathcal{A}, p) := \{a \in \mathcal{A} \mid p(a) \leq 1\}$ for the *closed ball semigroup* of $(\mathcal{A}, p)$.

Both $\mathtt{ball}(\mathcal{A}, p)$ and $\overline{\mathtt{ball}}(\mathcal{A}, p)$ are $*$-semigroups, and if $\mathcal{A}$ is unital, then $\mathbf{1} \in \overline{\mathtt{ball}}(\mathcal{A}, p)$. We will omit p from this notation whenever p is clear from the context, for instance if $\mathcal{A}$ is a C^*-algebra with the norm p. If $\mathcal{H}$ is a complex Hilbert space and $B(\mathcal{H})$ is the von Neumann algebra of all bounded linear operators on $\mathcal{H}$, then we also write $C(\mathcal{H}) := \overline{\mathtt{ball}}(B(\mathcal{H}))$.

(c) We define the *positive cone* of $\mathcal{A}$ as $\mathcal{A}_+ := \mathrm{conv}(\{a^*a \mid a \in \mathcal{A}\})$, where conv denotes the convex hull.

Here are some examples of seminormed involutive algebras.

Example 2.2. (a) Besides C^*-algebras and Banach $*$-algebras, a typical example for the above setting is the Fréchet algebra of smooth $\mathcal{A}_0$-valued functions $\mathcal{A} = \mathcal{C}^\infty(M, \mathcal{A}_0)$ on any compact manifold M, where $\mathcal{A}_0$ is any Banach $*$-algebra and $p(a) = \|a\|_\infty$. In this case $C^*(\mathcal{A}) = \mathcal{C}(M, \mathcal{A}_0)$ is the C^*-algebra of continuous $\mathcal{A}_0$-valued functions on M and $\eta_{\mathcal{A}}\colon \mathcal{C}^\infty(M, \mathcal{A}_0) \hookrightarrow \mathcal{C}(M, \mathcal{A}_0)$ is the isometric inclusion map.

(b) Let $(S, *, \alpha)$ be an involutive semigroup $(S, *)$, endowed with an *absolute value*, i.e., a non-negative function $\alpha\colon S \to [0,\infty)$ satisfying

$$\alpha(s^*) = \alpha(s) \quad \text{and} \quad \alpha(st) \leq \alpha(s)\alpha(t) \quad \text{for} \quad s, t \in S.$$

Then the real semigroup algebra $\mathcal{A} := \mathbb{R}[S]$ inherits an involution by the linear extension of the involution on S. Writing $\delta_s \in \mathcal{A}$ for the basis element corresponding to $s \in S$, we obtain by

$$p\Big(\sum_{s\in S} c_s\delta_s\Big) := \sum_{s\in S} |c_s|\alpha(s)$$

a seminormed real involutive algebra $(\mathbb{R}[S], p)$. (cf. [Ne00, Lemma III.2.4]).

Definition 2.3. (Enveloping C^*-algebras) For any seminormed involutive algebra $(\mathcal{A}, p)$, there exist a C^*-algebra $C^*(\mathcal{A}, p)$ and a morphism $\eta_{\mathcal{A}}\colon \mathcal{A} \to C^*(\mathcal{A}, p)$ of involutive algebras satisfying $\|\eta_{\mathcal{A}}(a)\| \leq p(a)$ with the universal property that, for every real linear homomorphism $\beta\colon \mathcal{A} \to \mathcal{B}$, where $\mathcal{B}$ is a C^*-algebra and $\|\beta(a)\| \leq p(a)$ for every $a \in \mathcal{A}$, there exists a unique morphism $\widetilde{\beta}\colon C^*(\mathcal{A}, p) \to \mathcal{B}$ of C^*-algebras satisfying $\widetilde{\beta} \circ \eta_{\mathcal{A}} = \beta$ (see [Ne00, Sect. III.2]). The morphism $\eta\colon \mathcal{A} \to C^*(\mathcal{A}, p)$ of real seminormed involutive algebras is called the *enveloping C^*-algebra of* $(\mathcal{A}, p)$.

Remark 2.4. (a) Once we have fixed the involutive seminorm p, it determines a topology on $\mathcal{A}$ so that mostly it is not necessary to specify any finer topology. However, for any topology τ on $\mathcal{A}$ which is finer than the one specified by p, the homomorphism $\eta_{\mathcal{A}}\colon \mathcal{A} \to C^*(\mathcal{A}, p)$ is continuous, so that every linear representation of $\mathcal{A}$ satisfying $\|\pi(a)\| \leq p(a)$ for $a \in \mathcal{A}$, is τ-continuous.

(b) A linear homomorphism $\pi\colon \mathcal{A} \to \mathcal{B}$, where $\mathcal{B}$ is a C^*-algebra satisfies $\|\pi(a)\| \leq p(a)$ for every $a \in \mathcal{A}$ if and only if $\pi(\mathtt{ball}(\mathcal{A}, p))$ is bounded (cf. Remark 3.5(b) below). This condition implies that $\|\pi(a)\| \leq q(a)$ holds for every equivalent submultiplicative involutive seminorm on $\mathcal{A}$. It follows in particular, that $C^*(\mathcal{A}, p) \cong C^*(\mathcal{A}, q)$ whenever there exist constants $c_1, c_2 > 0$ such that $c_1 q \leq p \leq c_2 q$.

(c) Since p is multiplicative on $\mathcal{A}$, the subspace $\mathcal{I}_p := \{a \in \mathcal{A}\colon p(a) = 0\}$ is an ideal of $\mathcal{A}$, and the quotient $\mathcal{A}/\mathcal{I}_p$ inherits the structure of a real normed involutive algebra. Accordingly, its completion is a real Banach $*$-algebra $\mathcal{A}_p$ whose enveloping C^*-algebra coincides with $C^*(\mathcal{A}, p)$.

We also point out that if $\mathcal{A}$ is a C^*-algebra, then $C^*(\mathcal{A})$ defined as above is not equal to $\mathcal{A}$. It is isomorphic to $\mathcal{A}_{\mathbb{C}} \simeq \mathcal{A} \oplus \overline{\mathcal{A}}$, since the universal property from Definition 2.3 involves all the real linear homomorphisms β, not only the complex linear ones.

Example 2.5. Suppose that $\mathcal{A}$ is commutative and let

$$\widehat{\mathcal{A}}_p := \{\chi \in \mathrm{Hom}(\mathcal{A}, \mathbb{R}) \setminus \{0\}\colon |\chi| \leq p\},$$

where $\mathrm{Hom}(\mathcal{A}, \mathbb{R})$ denotes the set of all $*$-homomorphisms $\mathcal{A} \to \mathbb{R}$. Note that every such homomorphism annihilates all skew-symmetric elements, so that we may assume that $a^* = a$ for every $a \in \mathcal{A}$. Then the weak-$*$-topology turns $\widehat{\mathcal{A}}_p$ into a locally compact space and it is easy to see that the natural map

$$\eta\colon \mathcal{A} \to C_0(\widehat{\mathcal{A}}_p) = C_0(\widehat{\mathcal{A}}_p, \mathbb{C}), \quad \eta(a) = \widehat{a}, \quad \widehat{a}(\chi) := \chi(a)$$

is the enveloping C^*-algebra of $(\mathcal{A}, p)$, so that $C^*(\mathcal{A}, p) \cong C_0(\widehat{\mathcal{A}}_p)$.

Example 2.6. (a) Consider $\mathcal{A} = M_n(\mathbb{K})$ for $\mathbb{K} \in \{\mathbb{R}, \mathbb{C}, \mathbb{H}\}$ with the natural involution $(a_{ij})^* = (\overline{a_{ji}})$ and any involutive seminorm p with $\|a\| \leq p(a)$ for $a \in \mathcal{A}$. Since $\mathcal{A}$ is simple, the ideal $\mathcal{I}_p := \{a \in \mathcal{A}\colon p(a) = 0\}$ is trivial, and since all non-degenerate representations of $\mathcal{A}$ are multiples of the complexification of the identical representations, we obtain

$$C^*(\mathcal{A}, p) \cong \mathcal{A}_{\mathbb{C}} \cong M_n(\mathbb{K}_{\mathbb{C}}) \cong \begin{cases} M_n(\mathbb{C}) & \text{for } \mathbb{K} = \mathbb{R} \\ M_n(\mathbb{C}) \oplus M_n(\mathbb{C}) & \text{for } \mathbb{K} = \mathbb{C} \\ M_n(M_2(\mathbb{C})) \cong M_{2n}(\mathbb{C}) & \text{for } \mathbb{K} = \mathbb{H}. \end{cases}$$

(b) For $\mathcal{A} := M_\infty(\mathbb{K}) := \bigcup_{n\in\mathbb{N}} M_n(\mathbb{K})$, one thus obtains for any $p \geq \|\cdot\|$ that

$$C^*(\mathcal{A},p) \cong \varinjlim C^*(M_n(\mathbb{K}),p) \cong K(\ell^2(\mathbb{N},\mathbb{K})_{\mathbb{C}}) \cong \begin{cases} K(\ell^2) & \text{for } \mathbb{K}=\mathbb{R} \\ K(\ell^2)\oplus K(\ell^2) & \text{for } \mathbb{K}=\mathbb{C} \\ K(\ell^2) & \text{for } \mathbb{K}=\mathbb{H}. \end{cases}$$

Completely positive functions

Definition 2.7. (a) A map $\varphi\colon \mathcal{A}\to\mathcal{B}$ between involutive algebras is said to be *positive* if $\varphi(\mathcal{A}_+)\subseteq\mathcal{B}_+$, i.e., positive elements of $\mathcal{A}$ are mapped to positive elements of $\mathcal{B}$. We call φ *completely positive* if, for every $n\in\mathbb{N}_0$, the induced map

$$\varphi_n\colon M_n(\mathcal{A})\to M_n(\mathcal{B}),\quad (a_{ij})\mapsto(\varphi(a_{ij}))$$

is positive, i.e., for $A=(a_{ij})\in M_n(\mathcal{A})_+$, the matrix $\varphi_n(A)\in M_n(\mathcal{B})$ is positive. This means that, for $B_1,\ldots,B_m\in M_n(\mathcal{A})$, we have $\varphi_n\big(\sum_{\ell=1}^m B_\ell^*B_\ell\big)\geq 0$. In view of

$$\Big(\sum_{\ell=1}^m B_\ell^*B_\ell\Big)_{ij} = \sum_{\ell,k} B^*_{\ell,ki}B_{\ell,kj},$$

this is equivalent to the requirement that

$$\Big(\varphi\Big(\sum_{k=1}^m a_{ki}^*a_{kj}\Big)\Big)_{1\leq i,j\leq n}\in M_n(\mathcal{B})_+ \quad \text{for} \quad (a_{ki})\in M_{m,n}(\mathcal{A}), m\in\mathbb{N}. \tag{1}$$

(b) This concept can be extended to maps defined on $\mathtt{ball}(\mathcal{A},p)$ as follows. We call a map $\varphi\colon \mathtt{ball}(\mathcal{A},p)\to\mathcal{B}$ *completely positive* if $A=(a_{ij})\in M_n(\mathcal{A})_+$ and $a_{ij}\in\mathtt{ball}(\mathcal{A},p)$ imply that the matrix $\varphi_n(A)=(\varphi(a_{ij}))\in M_n(\mathcal{B})$ is positive, for every $n\geq 1$.

Definition 2.8. Let S be a $*$-semigroup and $\mathcal{B}$ be a $*$-algebra. A function $\varphi\colon S\to\mathcal{B}$ is said to be *positive definite* if, for every $n\in\mathbb{N}$ and $s_1,\ldots,s_n\in S$, the matrix $(\varphi(s_j^*s_k))\in M_n(\mathcal{B})$ is positive.

The following lemma shows that completely positive functions are particular positive definite functions and one of the main points of the present paper is to explain in which sense they are particular among the positive definite functions.

Lemma 2.9. *If $\varphi\colon \mathtt{ball}(\mathcal{A},p)\to\mathcal{B}$ is completely positive, then φ is positive definite.*

Proof. If $a_1,\ldots,a_n\in\mathtt{ball}(\mathcal{A},p)$ and $A=(\delta_{1i}a_j)_{1\leq i,j\leq n}$, then $A^*A=(a_i^*a_j)_{1\leq i,j\leq n}$ with $p(a_i^*a_j)<1$. This implies the lemma. □

Remark 2.10. (a) If $\varphi\colon\mathcal{A}\to\mathcal{B}$ and $\psi\colon\mathcal{B}\to\mathcal{C}$ are completely positive maps, then so is their composition $\psi\circ\varphi\colon\mathcal{A}\to\mathcal{C}$. The same conclusion holds if $\varphi\colon\mathtt{ball}(\mathcal{A},p)\to\mathtt{ball}(\mathcal{B},q)\subseteq\mathcal{B}$ and $\psi\colon\mathtt{ball}(\mathcal{B},q)\to\mathcal{C}$ are completely positive, or if $\varphi\colon\mathtt{ball}(\mathcal{A},p)\to\mathcal{B}$ is completely positive and $\psi\colon\mathcal{B}\to\mathcal{C}$ is completely positive.

(b) If $\varphi\colon\mathcal{A}\to\mathcal{B}$ is a linear homomorphism of $*$-algebras, then so are the corresponding maps $\varphi_n\colon M_n(\mathcal{A})\to M_n(\mathcal{B})$, and this implies that φ is completely positive.

(c) From (1) it follows that a linear map $\varphi\colon \mathcal{A} \to \mathcal{B}$ is completely positive if and only if it is positive definite (cf. [Pau02, Lemma 3.13]; [Arv10, p. 112]). In particular, a linear functional $\varphi\colon \mathcal{A} \to \mathbb{C}$ is completely positive if and only if it is positive (cf. [St55, Thm. 3] for the case where $\mathcal{A}$ is a unital C^*-algebra).

(d) If $(S, *)$ is an involutive semigroup, then every map $\varphi\colon S \to \mathcal{B}$ extends to a linear map $\varphi_L\colon \mathbb{R}[S] \to \mathcal{B}$. Using (c), it is easy to see that φ_L is positive definite if and only if φ is positive definite.

The next lemma shows in particular that a homogeneous function on $\mathcal{A}$ is completely positive if and only if it is so on $\mathtt{ball}(\mathcal{A}, p)$.

Lemma 2.11. *Suppose that the map $\varphi\colon \mathtt{ball}(\mathcal{A}, p) \to \mathcal{B}$ is homogeneous of degree $\alpha \in \mathbb{R}$, i.e., for $0 < r < 1$ and $a \in \mathtt{ball}(\mathcal{A}, p)$, we have $\varphi(ra) = r^\alpha \varphi(a)$. Then φ extends by*

$$\widehat{\varphi}(ra) := r^\alpha \varphi(a) \quad \text{for} \quad 0 < r \text{ and } a \in \mathtt{ball}(\mathcal{A}, p)$$

to a map $\widehat{\varphi}\colon \mathcal{A} \to \mathcal{B}$. Then φ is completely positive if and only if $\widehat{\varphi}$ is completely positive.

Proof. We first observe that $\widehat{\varphi}$ is well-defined. It is clear that φ is completely positive if $\widehat{\varphi}$ is. To see the converse, we note that, for any matrix $A = (a_{ij}) \in M_n(\mathcal{A})_+$, there exists an $r > 0$ with $rp(a_{ij}) < 1$ for every i, j. Then $\widehat{\varphi}_n(A) = r^{-\alpha}\widehat{\varphi}_n(\alpha A) \in M_n(\mathcal{B})_+$ implies that $\widehat{\varphi}\colon \mathcal{A} \to \mathcal{B}$ is completely positive. □

Complexification of a real involutive algebra

Definition 2.12. (Complexification of a seminormed involutive algebra) Let $(\mathcal{A}, p)$ be a real seminormed involutive algebra. Then its complexification $\mathcal{A}_\mathbb{C}$ is an involutive algebra in the usual sense with respect to the antilinear extension $(x + iy)^* := x^* - iy^*$, for $x, y \in \mathcal{A}$, of the involution. Further,

$$p_\mathbb{C}(a + ib) := p(a) + p(b)$$

is a seminorm satisfying

$$p_\mathbb{C}((a + ib)^*) = p_\mathbb{C}(a^* - ib^*) = p(a^*) + p(b^*) = p(a) + p(b) = p_\mathbb{C}(a + ib)$$

and

$$\begin{aligned} p_\mathbb{C}((a + ib)(c + id)) &= p(ac - bd) + p(bc + ad) \le p(a)p(c) + p(b)p(d) + p(b)p(c) + p(a)p(d) \\ &= (p(a) + p(b))(p(c) + p(d)) = p_\mathbb{C}(a + ib)p_\mathbb{C}(c + id), \end{aligned}$$

so that $(\mathcal{A}_\mathbb{C}, p_\mathbb{C})$ is a seminormed involutive algebra. We call it the *complexification of* $(\mathcal{A}, p)$.

Lemma 2.13. *For a real involutive algebra $(\mathcal{A}, p)$, the complex linear extension*

$$\eta_\mathcal{A}^\mathbb{C}\colon \mathcal{A}_\mathbb{C} \to C^*(\mathcal{A}, p), \quad a + ib \mapsto \eta_\mathcal{A}(a) + i\eta_\mathcal{A}(b)$$

has the universal property of the enveloping C^*-algebra in the category of complex involutive algebras, i.e., for every complex linear $*$-homomorphism $\beta\colon \mathcal{A}_\mathbb{C} \to \mathcal{B}$ to a C^*-algebra satisfying $\|\beta(a)\| \le p_\mathbb{C}(a)$ for $a \in \mathcal{A}_\mathbb{C}$, there exists a unique morphism of C^*-algebras $\widehat{\beta}\colon C^*(\mathcal{A}, p) \to \mathcal{B}$ with $\widehat{\beta} \circ \eta_\mathcal{A}^\mathbb{C} = \beta$. In this sense we have $C^*(\mathcal{A}_\mathbb{C}, p_\mathbb{C}) \cong C^*(\mathcal{A}, p)$.

Proof. Let $\beta\colon \mathcal{A}_{\mathbb{C}} \to \mathcal{B}$ be a complex linear $*$-homomorphism to a C^*-algebra satisfying $\|\beta(a)\| \leq p_{\mathbb{C}}(a)$ for $a \in \mathcal{A}_{\mathbb{C}}$. Then $\|\beta(a)\| \leq p(a)$ for $a \in \mathcal{A}$, so that the universal property of $\eta_{\mathcal{A}}$ implies the existence of a morphism of C^*-algebras $\widehat{\beta}\colon C^*(\mathcal{A}, p) \to \mathcal{B}$ with $\widehat{\beta} \circ \eta_{\mathcal{A}} = \beta|_{\mathcal{A}}$. Now complex linear extension yields $\widehat{\beta} \circ \eta_{\mathcal{A}}^{\mathbb{C}} = \beta$. □

Remark 2.14. Note that a representation $\beta\colon \mathcal{A} \to B(\mathcal{H})$ satisfies $\|\beta(a)\| \leq p(a)$ for every $a \in \mathcal{A}$ if and only if its complex linear extension $\beta^{\mathbb{C}}$ satisfies $\|\beta^{\mathbb{C}}(a + ib)\| \leq p_{\mathbb{C}}(a + ib)$ for $a, b \in \mathcal{A}$.

Unitization of seminormed involutive algebras

Definition 2.15. Let $(\mathcal{A}, p)$ be a real seminormed involutive algebra and $\mathcal{A}^1 := \mathcal{A} \oplus \mathbb{R}$, $\mathbf{1} := (0, 1)$, the corresponding unital algebra with the involution $(a, \lambda)^* := (a^*, \lambda)$. Then $p^1(a, \lambda) := p(a) + |\lambda|$ turns $(\mathcal{A}^1, p^1)$ into a unital seminormed involutive algebra.

Remark 2.16. Every $*$-homomorphism $\beta\colon \mathcal{A} \to \mathcal{B}$ into a unital C^*-algebra $\mathcal{B}$, satisfying the condition $\|\beta(a)\| \leq p(a)$, extends by $\beta^1(a, \lambda) := \beta(a) + \lambda\mathbf{1}$ to a morphism of unital $*$-algebras satisfying $\|\beta^1(a)\| \leq p^1(a)$ for $a \in \mathcal{A}^1$. This implies that $C^*(\mathcal{A}^1, p^1)$ has the universal property of the *enveloping unital C^*-algebra* of $(\mathcal{A}, p)$, i.e.,

$$C^*(\mathcal{A}, p)^1 \cong C^*(\mathcal{A}^1, p^1),$$

(cf. [Bha98, Lemma 2.10] for the special case of Banach algebras).

Example 2.17. (a) For every $c \in \mathcal{A}$, the map $T_c\colon \mathcal{A}^1 \to \mathcal{A}, a \mapsto c^*ac$ is completely positive. For $n \in \mathbb{N}$, the induced map $T_{c,n}\colon M_n(\mathcal{A}^1) \to M_n(\mathcal{A})$ is given by $T_{c,n} = T_C$ with $C = cE_n$, where $E_n \in M_n(\mathcal{A})$ is the identity matrix.

For the sake of later reference, we note that this implies in particular that if $A = (a_{k\ell}) \in M_n(\mathcal{A}^1)_+ \subseteq M_n(M_N(\mathcal{A}^1)) \cong M_{nN}(\mathcal{A}^1)$ and $s_1, \ldots, s_N \in \mathcal{A}$, then the matrix $(s_i^* a_{k\ell} s_j) \in M_{nN}(\mathcal{A})$ is positive because it can be written as S^*AS for $S = (\delta_{ij} s_j)_{1 \leq i,j \leq N, 1 \leq k,\ell \leq n}$.

(b) Fix $s_1, \ldots, s_N \in \mathcal{A}$. We claim that the map

$$\beta\colon M_N(\mathcal{A}^1) \to \mathcal{A}, \quad \beta((a_{ij}) := \sum_{i,j=1}^N s_i^* a_{ij} s_j.$$

is completely positive. The corresponding map

$$\begin{aligned} \beta_n\colon M_n(M_N(\mathcal{A}^1)) \cong M_{nN}(\mathcal{A}^1) &\to M_n(\mathcal{A}), \\ \beta_n((a_{(i,k),(j,\ell)})) &= \Big(\sum_{i,j=1}^N s_i^* a_{(i,k),(j,\ell)} s_j \Big)_{1 \leq k,\ell \leq n} \end{aligned} \tag{2}$$

is of the same structure with s_i replaced by the matrix $s_i E_n \in M_n(\mathcal{A})$. Therefore it suffices to show that β is positive. As β is linear, this means that $\beta(B^*B) \geq 0$ for $B \in M_N(\mathcal{A}_1)$. For $A = B^*B$, we have

$$\beta(B^*B) = \sum_{i,k,j=1}^N s_i^* b_{ki}^* b_{kj} s_j = \sum_{k=1}^N \Big(\sum_{j=1}^N b_{kj} s_j \Big)^* \Big(\sum_{j=1}^N b_{kj} s_j \Big) \geq 0.$$

3 Positive definite functions and dilations

We start this section by a discussion of the representation $(\pi_\varphi, \mathcal{H}_\varphi)$ associated to a (bounded) positive definite function $\varphi\colon S \to B(V)$ on a $*$-semigroup S. Here $\mathcal{H}_\varphi$ is a subspace of the space V^S of V-valued functions on S, on which S acts by right translations. We then discuss dilations of positive definite functions and criteria for their existence, such as the existence of a unit in S. For non-unital semigroups, the existence of dilations is a subtle issue. Based on the concept of a structured $*$-semigroup and multiplier techniques, we prove in Theorem 3.18 the existence of dilations for positive definite functions on the canonical ideal S° of a structured $*$-semigroup, and this applies in particular to the subsemigroup $S^\circ = \mathtt{ball}(\mathcal{A}, p)$ of $S = \overline{\mathtt{ball}}(\mathcal{A}, p)$ when $\mathcal{A}$ is unital. Generalizing the classical Stinespring Dilation Theorem ([St55]), we prove in Proposition 3.9 that a linear completely positive function $\varphi\colon \mathcal{A} \to B(V)$ which is bounded on $\mathtt{ball}(\mathcal{A}, p)$ factors through $\eta_{\mathcal{A}}\colon \mathcal{A} \to C^*(\mathcal{A}, p)$ if and only if it has a dilation.

3.1 Preliminaries on positive definite functions

Throughout this subsection, $(S, *)$ is an involutive semigroup.

Remark 3.1. (The GNS representation associated to φ) We recall from [Ne00] some basic facts on the reproducing kernel Hilbert space associated with any positive definite function $\varphi\colon S \to B(V)$, where S is any $*$-semigroup. Define the positive definite kernel

$$K = K_\varphi\colon S \times S \to B(V) \quad \text{by} \quad K(s,t) := \varphi(st^*).$$

The corresponding reproducing kernel space $\mathcal{H}_\varphi \subseteq V^S$ (V^S stands for the vector space of all V-valued functions on S), can be constructed as follows. For $t \in S, v \in V$, we put $K_{t,v} := K(\cdot, t)v \in V^S$. Then

$$\mathcal{H}_\varphi^0 := \operatorname{span}\{K_{t,v} \mid t \in S,\ v \in V\}$$

is a pre-Hilbert space with respect to a scalar product specified by

$$\langle K_{s,v}, K_{t,w}\rangle = \langle K(t,s)v, w\rangle_V = \langle \varphi(ts^*)v, w\rangle_V \quad \text{for} \quad s,t \in S, v, w \in V.$$

The completion $\mathcal{H}_\varphi$ of $\mathcal{H}_\varphi^0$ has a natural inclusion into V^S with continuous evaluation operators

$$K_s\colon \mathcal{H}_\varphi \to V, f \mapsto f(s) \qquad \text{satisfying} \quad K_s K_t^* = K(s,t) \quad \text{for} \quad s,t \in S.$$

In particular, we have $K_{t,v} = K_t^* v$ for $t \in S, v \in V$.

As in [Ne00, Sect. II.3], we denote by $B^0(\mathcal{H}_\varphi^0)$ the $*$-algebra consisting of the linear operators $A\colon \mathcal{H}_\varphi^0 \to \mathcal{H}_\varphi^0$ for which there exists a linear operator $A^\sharp\colon \mathcal{H}_\varphi^0 \to \mathcal{H}_\varphi^0$ with $\langle Av, w\rangle = \langle v, A^\sharp w\rangle$ for all $v, w \in \mathcal{H}_\varphi^0$. Then we obtain a $*$-representation $\pi_\varphi^0\colon S \to B^0(\mathcal{H}_\varphi^0)$ given by $(\pi_\varphi^0(s)f)(t) := f(ts)$ for all $t, s \in S$. For the operator closures $\pi_\varphi(s) := \overline{\pi_\varphi^0(s)}$, we then obtain

$$K_t \pi_\varphi(s) = K_{ts} \quad \text{and} \quad \pi_\varphi(s) K_t^* = K_{ts^*} \quad \text{for} \quad t, s \in S. \tag{3}$$

Remark 3.2. If $\varphi\colon S \to B(V)$ is constant, then all functions $K_{t,v}(s) = K(s,t)v = \varphi(st^*)v$ are constant, so that the representation $(\pi_\varphi, \mathcal{H}_\varphi)$ is trivial.

If, conversely, the representation π_φ is trivial, then $K_t = K_{ts}$ for $t, s \in S$ by (3). If S has a unit, this implies that φ is constant.

Definition 3.3. Let S be a $*$-semigroup, V be a complex Hilbert space and $\varphi\colon S \to B(V)$ be a positive definite function.

(a) If $\alpha\colon S \to [0,\infty)$ is an absolute value (cf. Examples 2.2(b)), then φ is said to be *α-bounded* if

$$\varphi(s^*t^*ts) \leq \alpha(t)^2\varphi(s^*s) \quad \text{for} \quad t, s \in S.$$

This condition is equivalent to $\|\pi_\varphi(t)\| \leq \alpha(t)$ for the corresponding GNS representation $(\pi_\varphi, \mathcal{H}_\varphi)$ (cf. [Ne00, Th. III.1.3]; and [Seb86] for related conditions).

(b) A triple $(\pi, \mathcal{H}, \iota)$ consisting of a $*$-representation $\pi\colon S \to B(\mathcal{H})$ on the complex Hilbert space $\mathcal{H}$ and a bounded linear operator $\iota\colon V \to \mathcal{H}$ (called the *linking operator*), is called a *dilation of φ* if

$$\varphi(s) = \iota^*\pi(s)\iota \quad \text{for} \quad s \in S.$$

A dilation is said to be *minimal* if $\pi(S)\iota(V)$ spans a dense subspace of $\mathcal{H}$. A positive definite function for which a dilation exists is called *dilatable*. [1]

The following proposition shows in particular that the boundedness characterization of [Ne00, Th. III.1.19(4)] can also be used for operator-valued functions.

Proposition 3.4. *For any positive definite function $\varphi\colon S \to B(V)$ on the involutive semigroup $(S, *)$, the following assertions hold:*

(a) *If S is unital and φ is α-bounded for some absolute value α on S, then $(\pi_\varphi, \mathcal{H}_\varphi, K_{\mathbf{1}}^*)$ is a minimal dilation of φ satisfying $\|\pi_\varphi(s)\| \leq \alpha(s)$ for $s \in S$.*

(b) *If $(\pi, \mathcal{H}, \iota)$ is a minimal dilation of φ, then there exists a unique unitary operator $\Phi\colon \mathcal{H} \to \mathcal{H}_\varphi$ intertwining π with π_φ and satisfying $\Phi(\iota(v)) = \varphi(\cdot)\cdot v$ for all $v \in V$.*

(c) (Uniqueness of minimal dilations) *If $(\pi_j, \mathcal{H}_j, \iota_j)$, $j = 1, 2$, are two minimal dilations of the positive definite function $\varphi\colon S \to B(V)$, then there exists a unique unitary operator $U\colon \mathcal{H}_1 \to \mathcal{H}_2$ intertwining π_1 with π_2 and satisfying $U \circ \iota_1 = \iota_2$.*

Proof. (a) We will use the notation from Remark 3.1. In view of [Ne00, Th. III.1.3], the α-boundedness of φ implies that $\|\pi_\varphi^0(s)\| \leq \alpha(s)$ for every $s \in S$. Since $\mathcal{H}_\varphi^0$ is dense in $\mathcal{H}_\varphi$, it follows that the linear operator $\pi_\varphi^0(s) \in B(\mathcal{H}_\varphi^0)$ has a unique extension $\pi_\varphi(s) \in B(\mathcal{H}_\varphi)$ with $\|\pi_\varphi(s)\| \leq \alpha(s)$. This is also given by right translations

$$(\pi_\varphi(s)f)(t) := f(ts) \quad \text{for} \quad t, s \in S, f \in \mathcal{H}_\varphi$$

(cf. [Ne00, Prop. II.4.9]). For $s \in S$ and $\iota := K_{\mathbf{1}}^*$, we have

$$\iota^*\pi_\varphi(s)\iota(v) = K_{\mathbf{1}}\pi_\varphi(s)K_{\mathbf{1}}^*v = (K_{\mathbf{1}}^*v)(s) = \varphi(s)v,$$

[1] The above dilatable functions were called Stinespring representable in [Bha98] in the special case when $S = (\mathcal{A}, \cdot)$ for some Banach $*$-algebra $\mathcal{A}$.

so that $(\pi_\varphi, \mathcal{H}_\varphi, \iota)$ is a dilation of φ. Its minimality follows from the fact that, for $f \in \mathcal{H}_\varphi$, the vanishing of $f(s) = K_{\mathbf{1}}\pi(s)f$ for every $s \in S$ implies $f = 0$.

(b) We consider the map

$$\Phi\colon \mathcal{H} \to V^S, \quad \Phi(w)(s) := \iota^*\pi(s)w.$$

This map is S-equivariant with respect to the right translation action of S on V^S and satisfies $\Phi(\iota(v)) = \varphi \cdot v$ for all $v \in V$. If $\Phi(w) = 0$, then $\pi(S)w \subseteq \ker \iota^* = \iota(V)^\perp$ leads to $w \in (\pi(S)\iota(V))^\perp = \{0\}$, so that Φ is injective. Therefore $\Phi(\mathcal{H}) \subseteq V^S$ is a reproducing kernel Hilbert space whose kernel is given by

$$K(s,t) = (\iota^*\pi(s))(\iota^*\pi(t))^* = \iota^*\pi(st^*)\iota = \varphi(st^*).$$

We conclude that $(\pi, \mathcal{H})$ is equivalent to $(\pi_\varphi, \mathcal{H}_\varphi)$.

(c) follows from (b) with $U := \Phi_2^{-1} \circ \Phi_1$. □

Remark 3.5. Let S be a $*$-semigroup, $\varphi\colon S \to B(V)$ be a positive definite function and $\alpha\colon S \to [0,\infty)$ be an absolute value.

(a) If $\|\varphi(s)\| \leq M\alpha(s)$ for all $s \in S$ and some $M \geq 0$, then φ is α-bounded (see the supplement to [RSzN72, Appendix 1] or [Ne00, Cor. III.1.20]). In particular, Proposition 3.4(a) applies to every bounded positive definite function $\varphi\colon S \to B(V)$ with the constant absolute value $\alpha = 1$. [2]

(b) If $\pi\colon S \to B(\mathcal{H})$ is a $*$-representation for which $\sup\limits_{s\in S}\|\pi(s)\| < \infty$, then actually $\sup\limits_{s\in S}\|\pi(s)\| \leq 1$ because, for $s \in S$, the boundedness of the sequence

$$\|\pi(s)\|^{2n} = \|\pi(s^*s)\|^n = \|\pi((s^*s)^n)\|$$

is equivalent to $\|\pi(s)\| \leq 1$.

Lemma 3.6. *Let S be a $*$-semigroup with a zero element 0, i.e., $0 \cdot s = s \cdot 0 = 0$ for $s \in S$. If $\varphi\colon S \to B(V)$ is an α-bounded positive definite function, then the function $\varphi - \varphi(0)$ is also positive definite.*

Proof. Since 0 is a zero element, the same holds for 0^*, and this leads to $0 = 0^*0 = 0^*$. Therefore the operator $P := \pi_\varphi(0)$ is a hermitian projection satisfying $P\pi_\varphi(s) = \pi_\varphi(s)P = P$ for every $s \in S$. It acts on $\mathcal{H}_\varphi$ by

$$(\pi_\varphi(0)f)(s) = f(s0) = f(0).$$

This implies that $\mathcal{H}_0 := P(\mathcal{H}_\varphi)$ is the subspace of constant functions contained in $\mathcal{H}_\varphi$, and that $\mathcal{H}_0^\perp = \{f \in \mathcal{H}_\varphi\colon f(0) = 0\}$. Accordingly, the reproducing kernel $K(s,t) = \varphi(st^*)$ decomposes as

$$K_s = \mathrm{ev}_0 \oplus (K_s - \mathrm{ev}_0) \in B(\mathcal{H}_0, V) \oplus B(\mathcal{H}_0^\perp, V) \cong B(\mathcal{H}_\varphi, V).$$

This leads to

$$\begin{aligned}(K_s - \mathrm{ev}_0)(K_t - \mathrm{ev}_0)^* &= K_sK_t^* - \mathrm{ev}_0\, K_t^* - K_s\, \mathrm{ev}_0^* + \mathrm{ev}_0\, \mathrm{ev}_0^* \\ &= \varphi(st^*) - \varphi(0) - \varphi(0) + \varphi(0) = \varphi(st^*) - \varphi(0).\end{aligned}$$

We conclude that $\varphi - \varphi(0)$ is positive definite. □

[2] In the case $V = \mathbb{C}$ one thus recovers [Arv87, Lemma 2.3].

Proposition 3.7. (Existence of minimal dilations) *Consider the unital involutive semigroup $S^1 := S \dot\cup \{\mathbf{1}\}$ with $\mathbf{1}^* = \mathbf{1}$ and an absolute value α on S^1. For an α-bounded positive definite function $\varphi\colon S \to B(V)$, the following are equivalent:*

(i) *φ is dilatable.*

(ii) *For every $v \in V$, the function $\varphi \cdot v\colon S \to V$ belongs to $\mathcal{H}_\varphi$.*

(iii) *φ extends to an α-bounded positive definite function on S^1.*

Proof. (i) $\Rightarrow$ (ii) follows from Proposition 3.4(b).

(ii) $\Rightarrow$ (i): Suppose, conversely, that $\varphi \cdot v \in \mathcal{H}_\varphi$ for every $v \in V$. Then we obtain a linear map $\iota\colon V \to \mathcal{H}_\varphi, v \mapsto \varphi \cdot v$. Since all compositions $\mathrm{ev}_s \circ \iota\colon V \to V, v \mapsto \varphi(s)v$, are continuous, the Closed Graph Theorem implies that ι is continuous. We further have, for $v, w \in V$, the relation

$$\langle \iota^* \pi_\varphi(s^*)\iota(v), w\rangle = \langle \pi_\varphi(s^*)\varphi \cdot v, \varphi \cdot w\rangle = \langle K_s^* v, \varphi \cdot w\rangle = \langle v, \varphi(s)w\rangle.$$

This implies that $\varphi(s) = (\iota^* \pi_\varphi(s^*)\iota)^* = \iota^* \pi_\varphi(s)\iota$, so that $(\pi_\varphi, \mathcal{H}_\varphi, \iota)$ is a dilation of φ.

(i) $\Rightarrow$ (iii): If $(\pi, \mathcal{H}, \iota)$ is a dilation of φ and $\pi^1(\mathbf{1}) := \mathbf{1}$ is the canonical extension of π to a representation of S^1, then $\varphi^1(s) := \iota^* \pi^1(s)\iota$ is an extension of φ to an α-bounded positive definite function on S^1.

(iii) $\Rightarrow$ (i): If an α-bounded positive definite extension φ^1 to S^1 exists, then Proposition 3.4(a) implies that φ^1, and hence also φ, is dilatable. □

Example 3.8. (a) If $\varphi\colon S \to B(V)$ is a $*$-representation, then $(\varphi, V, \mathrm{id}_V)$ is a dilation of φ.

(b) If $\varphi\colon S \to B(V)$ is positive definite and $s \in S$, then $\varphi_s(t) := \varphi(sts^*)$ defines a positive definite function on S which extends by $\varphi_s(\mathbf{1}) := \varphi(ss^*)$ to a positive definite function on $S^{\mathbf{1}}$. Therefore φ_s is dilatable by Proposition 3.7. More explicitly, the representation $(\pi_\varphi, \mathcal{H}_\varphi)$ satisfies

$$K_s \pi_\varphi(t) K_s^* = K_{st} K_s^* = \varphi(sts^*) = \varphi_s(t),$$

so that $(\pi_\varphi, \mathcal{H}_\varphi, K_s^*)$ is a dilation of φ_s.

(c) Suppose that S is endowed with a topology for which the bounded positive definite function $\varphi\colon S \to B(V)$ is weak-operator continuous and for which there exists a right approximate identity $(\delta_j)_{j\in J}$ in S, i.e., $\lim_j s\delta_j = s$ for every $s \in S$.

For $v \in V$, we then have $\|K_{\delta_j, v}\|^2 = \langle \varphi(\delta_j \delta_j^*)v, v\rangle \le \|\varphi\|_\infty \|v\|^2$, so that the net $(K_{\delta_j,v})_{j\in J}$ has a weak cluster point $\psi \in \mathcal{H}_\varphi$. Let $K_{\delta_{j_k}, v}$ be a subnet converging to ψ. Then we obtain for $w \in V$ the relation

$$\langle \psi(s), w\rangle = \lim_k \langle K_{\delta_{j_k}, v}, K_{s,w}\rangle = \lim_k \langle \varphi(s\delta_{j_k})v, w\rangle = \langle \varphi(s)v, w\rangle,$$

so that $\psi = \varphi \cdot v \in \mathcal{H}_\varphi$. Therefore φ is dilatable by Proposition 3.7(iii).

Proposition 3.9. *Let $(\mathcal{A}, p)$ be a seminormed $*$-algebra and V be a complex Hilbert space. Then $\Phi \mapsto \Phi \circ \eta_\mathcal{A}$ is a bijective correspondence between the set of linear completely positive $B(V)$-valued maps on $C^*(\mathcal{A}, p)$ and the dilatable completely positive functions $\varphi\colon \mathcal{A} \to B(V)$ which are bounded on $\mathtt{ball}(\mathcal{A}, p)$.*

Proof. Since $\eta_{\mathcal{A}}$ is a linear homomorphism of $*$-algebras, it is completely positive (Remark 2.10) and its range spans a dense subspace of $C^*(\mathcal{A},p)$. Hence the correspondence $\Phi \mapsto \Phi\circ\eta_{\mathcal{A}}$ is well defined and injective. Every completely positive map $\Phi\colon C^*(\mathcal{A},p)\to B(V)$ extends to a completely positive map $\Phi^1\colon C^*(\mathcal{A},p)^1\to B(V)$ by Arveson's Extension Theorem ([Arv69, Thm. 1.2.3]), and $C^*(\mathcal{A},p)^1\cong C^*(\mathcal{A}^1,p^1)$ by Remark 2.16. Therefore Φ is dilatable, and this implies that $\Phi\circ\eta_{\mathcal{A}}$ is dilatable.

Assume, conversely, that $\varphi\colon \mathcal{A}\to B(V)$ is linear, bounded on $\mathtt{ball}(\mathcal{A},p)$, and that $(\pi,\mathcal{H},\iota)$ is a minimal dilation of φ. Then the linearity of φ implies that the representation π of $(\mathcal{A},\cdot)$ is linear. Further, $\|\pi(a)\|\le p(a)$ follows from the uniqueness of minimal dilations (Proposition 3.4). By the universal property of $C^*(\mathcal{A},p)$, there exists a unique $*$-homomorphism $\widehat{\pi}\colon C^*(\mathcal{A},p)\to B(\mathcal{H})$ with $\widehat{\pi}\circ\eta_{\mathcal{A}}=\pi$. Then $\Phi(\cdot):=\iota^*\widehat{\pi}(\cdot)\iota\colon C^*(\mathcal{A},p)\to B(V)$ is linear, completely positive and satisfies $\Phi\circ\eta_{\mathcal{A}}=\varphi$. This completes the proof. □

A Banach $*$-algebra version of Proposition 3.9 can be found in [Bha98, Th. 2.1(6)]. In that case the $*$-representations and linear positive definite $B(V)$-valued functions are always continuous ([Dix64, Prop. 1.3.7], [Arv10, Cor. A.2]). This automatic continuity property remains true for unital Mackey complete $*$-algebras with continuous inversion ([Bi10, Prop. 6.6]).

Example 3.10. (A non-dilatable completely positive functional) We consider the non-unital Banach $*$ algebra $\mathcal{A}:=\ell^1(\mathbb{N},\mathbb{R})$, endowed with the pointwise multiplication and $p(a)=\|a\|_1$. Then the enveloping C^*-algebra is $C^*(\mathcal{A},p)\cong c_0(\mathbb{N},\mathbb{C})$.

The continuous linear functional

$$\varphi\colon \mathcal{A}\to\mathbb{R},\quad \varphi(a):=\sum_{n=1}^{\infty}a_n$$

is a sum of characters of $\mathcal{A}$, hence completely positive. Moreover, for $a,b\in\mathcal{A}$, we have

$$\varphi(a^*ba)\le\|b\|_\infty\varphi(a^*a)=\|b\|_\infty\|a\|_2^2,$$

so that the representation π_φ of $\mathcal{A}$ on $\mathcal{H}_\varphi\subseteq\mathcal{A}^*=\ell^\infty(\mathbb{N},\mathbb{C})$ extends to a representation of $C^*(\mathcal{A},p)\cong c_0(\mathbb{N},\mathbb{C})$ (Proposition 3.4).

Note that $\varphi(ab^*)=\sum_n a_n\overline{b_n}$ implies that $\mathcal{H}_\varphi\cong\ell^2(\mathbb{N},\mathbb{C})$, considered as a subspace of $\mathcal{A}'_{\mathbb{C}}\cong\ell^\infty(\mathbb{N},\mathbb{C})$. The function φ itself corresponds to the constant function 1, which is not contained in $\mathcal{H}_\varphi$. This implies that there exists no element $v\in\mathcal{H}_\varphi$ with $\varphi(a)=\langle\pi(a)v,v\rangle$ for $a\in\mathcal{A}$ (cf. [Ne00, Cor. III.1.25]). We conclude that φ is not dilatable (Proposition 3.7).

Note that $\mathcal{A}$ has no bounded approximate identity because, in view of Example 3.8(c), this would contradict Proposition 3.7, applied to $\mathtt{ball}(\mathcal{A},p)$ (see also [Ne00, Rem. IV.1.23]).

The preceding example, applied to $\mathtt{ball}(\mathcal{A},p)$, provides a counterexample to [Seb86, Th. 2].

3.2 Multipliers and structured $*$-semigroups

In this subsection we recall the multiplier semigroup of a $*$-semigroup and introduce the concept of a structured (unital) $*$-semigroup (S,γ), which consists of a $*$-semigroup S with a unital $*$-homomorphism $\gamma\colon(0,1]\to S$ with central range. The main result of this subsection is Theorem 3.18 which asserts that bounded positive definite functions on the semigroup ideal $S^\circ:=\gamma((0,1))S$ are always dilatable.

Multipliers of involutive semigroups

Definition 3.11. (Multiplier semigroup) Let $(S, *)$ be an involutive semigroup. A *multiplier* of S is a pair (λ, ρ) of mappings $\lambda, \rho\colon S \to S$ satisfying the following conditions:

(M1) $a\lambda(b) = \rho(a)b$,

(M2) $\lambda(ab) = \lambda(a)b$, and

(M3) $\rho(ab) = a\rho(b)$.

The map λ is called the *left action* of the multiplier and ρ is called the *right action* of the multiplier. For $m = (\lambda, \rho)$ and $s \in S$ we write

$$ms := \lambda(s) \quad \text{and} \quad sm := \rho(s).$$

We write $M(S)$ for the set of all multipliers of S and turn it into an involutive semigroup by

$$(\lambda, \rho)(\lambda', \rho') := (\lambda \circ \lambda', \rho' \circ \rho) \quad \text{and} \quad (\lambda, \rho)^* := (\rho^*, \lambda^*)$$

with $\lambda^*(a) := \lambda(a^*)^*$ and $\rho^*(a) = \rho(a^*)^*$ (see [Jo64]).

Remark 3.12. (a) The assignment $\eta\colon S \to M(S), a \mapsto (\lambda_a, \rho_a)$ defines a morphism of involutive semigroups which is surjective if S has an identity: in this case (M1) implies that $s := \lambda(\mathbf{1}) = \rho(\mathbf{1})$, (M2) implies $\lambda = \lambda_s$ and (M3) that $\rho = \rho_s$. The multiplier semigroup acts from the right on S by $(s, m) \mapsto sm$, and via η this extends the natural right action of S on itself. This right action of $M(S)$ is related to the fact that $\eta(S)$ is an involutive semigroup ideal in $M(S)$. In fact, $\eta(s)^* = (\rho_s^*, \lambda_s^*) = (\lambda_{s^*}, \rho_{s^*}) = \eta(s^*)$, and

$$(\lambda, \rho)\eta(s) = (\lambda \circ \lambda_s, \rho_s \circ \rho) = (\lambda_{\lambda(s)}, \rho_{\lambda(s)}) = \eta\big(\lambda(s)\big).$$

(b) Let φ be a positive definite function on S and $K_\varphi(a, b) := \varphi(ab^*)$ the corresponding positive definite kernel. We claim that K_φ is invariant with respect to the action of $M(S)$ on S, as is easily seen from

$$K_\varphi(a, bm) = \varphi(a(bm)^*) = \varphi(am^*b^*) = K_\varphi(am^*, b).$$

This shows that, for every positive definite function φ on S, there exists a hermitian representation $\widetilde{\pi}_\varphi$ of $M(S)$ on $\mathcal{H}_\varphi^0$ satisfying $\widetilde{\pi}_\varphi \circ \eta = \pi_\varphi$. In view of [Ne00, Prop. II.2.11(ii)], every bounded representation $(\pi, \mathcal{H})$ of S yields a hermitian representation $\widetilde{\pi}$ of the multiplier semigroup with $\widetilde{\pi}(m)\pi(s) = \pi(ms)$ and therefore also $\widetilde{\pi} \circ \eta = \pi$. To decide whether this extension acts by bounded operators, one needs more information on the multipliers or has to use topological arguments involving approximate identities (cf. [FD88, p.778]).

Lemma 3.13. *For any non-degenerate bounded representation $(\pi, \mathcal{H})$ of the involutive semigroup $(S, *)$, there exists a unique bounded representation $(\widetilde{\pi}, \mathcal{H})$ of its multiplier semigroup $M(S)$ satisfying $\widetilde{\pi}(m)\pi(s) = \pi(ms)$ for $m \in M(S), s \in S$.*

Proof. Let $(\pi, \mathcal{H})$ be a non-degenerate bounded representation of S and note that this implies that $\|\pi(s)\| \leq 1$ for every $s \in S$. We claim that the corresponding hermitian representation $\widetilde{\pi}$ of $M(S)$ is also bounded. To see this, we may w.l.o.g. assume that π is cyclic. So let $v \in \mathcal{H}$ be an element for which $\pi(S)v$ is total in $\mathcal{H}$. Then $\widetilde{\pi}(M(S))\pi(S)v \subseteq \pi(S)v$ shows that this total subset is invariant under the hermitian representation of $M(S)$ on $\mathrm{span}(\pi(S)v)$. From $\|\pi(s)v\| \leq \|v\|$ it now follows that, for every $s \in S$, the positive definite function

$$\varphi(m) := \langle \widetilde{\pi}(m)\pi(s)v, \pi(s)v\rangle = \langle \pi(s^*ms)v, v\rangle$$

is bounded, so that [Ne00, Cor. III.1.20] implies that

$$\|\widetilde{\pi}(m)\pi(s)v\| \leq \|\pi(s)v\|.$$

Therefore [Ne00, Lemma II.3.8(iii)] yields $\|\widetilde{\pi}(m)\| \leq 1$ for every multiplier $m \in M(S)$. □

Example 3.14. Let $(\mathcal{A}, p)$ be a real seminormed involutive algebra and $(\mathcal{A}^1, p^1)$ be the corresponding unital seminormed algebra (Definition 2.15). Then $\mathcal{A}$ is an ideal in $\mathcal{A}^1$ and $\mathtt{ball}(\mathcal{A}, p)$ is a semigroup ideal in $\overline{\mathtt{ball}}(\mathcal{A}^1, p^1)$, which leads to a natural homomorphism

$$\mu\colon \overline{\mathtt{ball}}(\mathcal{A}^1, p^1) \to M(\mathtt{ball}(\mathcal{A}, p)), \quad \mu(m) := (\lambda_m, \rho_m).$$

Accordingly, Lemma 3.13 implies that every non-degenerate bounded representation π of $\mathtt{ball}(\mathcal{A}, p)$ extends uniquely to a representation π^1 of $\overline{\mathtt{ball}}(\mathcal{A}^1, p^1)$ satisfying $\pi^1(m)\pi(a) = \pi(ma)$ for $a \in \mathtt{ball}(\mathcal{A}, p)$, $m \in \overline{\mathtt{ball}}(\mathcal{A}^1, p^1)$.

Structured $*$-semigroups

Definition 3.15. A *structured $*$-semigroup* is a unital $*$-semigroup S with a homomorphism of $*$-semigroups $\gamma\colon (0,1] \to S$ for which $\gamma(1) = \mathbf{1} \in S$ and the image of γ is contained in the center of S. We will say that γ is the *structure homomorphism* of S. Note that $\gamma(r)^* = \gamma(r)$ for every $r \in (0,1]$. We will denote

$$(\forall r \in (0,1])(\forall x \in S) \quad xr = rx := \gamma(r)x$$

and $S^\circ := \bigcup_{0<r<1} rS$, which is easily seen to be a $*$-subsemigroup of S, however we may have $\mathbf{1} \notin S^\circ$. Actually, S° is a semigroup ideal of S, that is, $SS^\circ \cup S^\circ S \subseteq S^\circ$.

Remark 3.16. If S is a structured $*$-semigroup, then $S^\circ = S^\circ S^\circ$, as a trivial consequence of $S \cdot S = S$ (S is unital).

Every $*$-semigroup with unit element is a structured $*$-semigroup with respect to the trivial structure homomorphism which is identically equal to the unit element. Here are also some less trivial examples.

Example 3.17. (a) Let $\mathcal{A}$ be any unital associative $*$-algebra, and pick any convex self-adjoint multiplicative subset $S \subseteq \mathcal{A}$ which contains $0, \mathbf{1} \in \mathcal{A}$. That is, we require that S should be convex, closed under the involution, and moreover $S \cdot S \subseteq S$ and $0, \mathbf{1} \in S$. Then the multiplicative $*$-semigroup S is a typical example of a structured $*$-semigroup with the structure homomorphism defined by the scalar multiplication $\gamma\colon (0,1] \to S$, $\gamma(r) = r\mathbf{1}$.

(b) Let $(\mathcal{A}, p)$ is any seminormed unital $*$-algebra. Then the $*$-semigroup

$$S := \overline{\mathtt{ball}}(\mathcal{A}, p) := \{a \in \mathcal{A} \colon p(a) \leq 1\}$$

satisfies the requirements of (a). It is easily checked that

$$S^\circ = \mathtt{ball}(\mathcal{A}, p) := \{a \in \mathcal{A} \colon p(a) < 1\}.$$

(c) Let $\mathcal{A}$ be a unital associative Banach $*$-algebra with isometric involution and $\|\mathbf{1}\| = 1$. Then the contraction $*$-semigroup $S = \overline{\mathtt{ball}}(\mathcal{A}) = \{a \in \mathcal{A} \colon \|a\| \leq 1\}$ is a special case of (b) and it is clear that $S^\circ = \mathtt{ball}(\mathcal{A})$ is the strict contraction semigroup of $\mathcal{A}$.

We will prove the following generalization of [Arv87, Th. 2.2]. Note that it does not follows from Proposition 3.4 because S° need not be unital.

Theorem 3.18. (Dilation Theorem for structured $*$-semigroups) *Let S be a structured $*$-semigroup and V be a complex Hilbert space. Then every bounded positive definite function $\varphi \colon S^\circ \to B(V)$ has a dilation $(\pi, \mathcal{H}, \iota)$ with*

$$\|\iota\|^2 \leq \limsup_{0<r<1} \|\varphi(\gamma(r))\|.$$

Before proving Theorem 3.18, we describe some of its consequences.

Remark 3.19. (a) If $(\mathcal{A}, p)$ is a unital seminormed $*$-algebra, then $S = \overline{\mathtt{ball}}(\mathcal{A}, p)$ is a structured $*$-semigroup with $S^\circ = \mathtt{ball}(\mathcal{A}, p)$ (Example 3.17(c)). Accordingly, Theorem 3.18 applies to S.

(b) Example 3.10 shows that Theorem 3.18 does not generalize to non-unital semigroups.

For later use in the proof of Proposition 3.21, we also record the following observation.

Lemma 3.20. *If S is a $*$-semigroup satisfying $S = S \cdot S$, and $\varphi \colon S \to B(V)$ is an α-bounded positive definite function, then the representation $(\pi_\varphi, \mathcal{H}_\varphi)$ is non-degenerate.*

Proof. Let $f \in \mathcal{H}_\varphi$ with $\pi_\varphi(S)f = \{0\}$. Then $0 = f(S \cdot S) = f(S)$ implies $f = 0$. □

We now turn to the proof of Theorem 3.18, which needs the following method of extending representations of semigroup ideals. In the next proposition we use the notation S° from structured $*$-semigroups only in order to indicate how this method will be used, however this applies to semigroup $*$-ideals, which are more general than the ones defined by a structure homomorphism as in Definition 3.15.

Proposition 3.21. *Let S be a $*$-semigroup and suppose that $S^\circ \trianglelefteq S$ is a semigroup $*$-ideal, i.e., $SS^\circ \cup S^\circ S \subseteq S^\circ$, and that it satisfies $S^\circ S^\circ = S^\circ$. Let $\varphi \colon S^\circ \to B(V)$ be a bounded positive definite function with the corresponding representation $\pi_\varphi \colon S^\circ \to B(\mathcal{H}_\varphi)$ on the reproducing kernel Hilbert space $\mathcal{H}_\varphi$ consisting of V-valued functions on S°. Then the formula*

$$(\widehat{\pi}_\varphi(s)f)(t) = f(ts) \tag{4}$$

defines a representation $\widehat{\pi}_\varphi \colon S \to B(\mathcal{H}_\varphi)$ by contractions, which extends the representation π_φ.

Proof. Since $\varphi\colon S^\circ \to B(V)$ is a bounded positive definite function, we obtain the corresponding GNS representation $(\pi_\varphi, \mathcal{H}_\varphi)$ of S°, where $\mathcal{H}_\varphi \subseteq V^{S^\circ}$. The assumption $S^\circ S^\circ = S^\circ$ implies that this representation is non-degenerate (Lemma 3.20).

Since S acts in the obvious way by multipliers on the semigroup ideal S°, Lemma 3.13 implies the existence of a bounded representation $\widehat{\pi}_\varphi$ of S on $\mathcal{H}_\varphi$ which is uniquely determined by $\widehat{\pi}_\varphi(s)\pi_\varphi(t) = \pi_\varphi(st)$ for $s \in S$, $t \in S^\circ$. To see that this representation is also given by (4), it suffices to verify this on functions of the form $f = \pi_\varphi(t)h$, $h \in \mathcal{H}_\varphi$:

$$(\widehat{\pi}_\varphi(s)\pi_\varphi(t)h)(x) = (\pi_\varphi(st)h)(x) = h(xst) = (\pi_\varphi(t)h)(xs).$$

This completes the proof. □

The following lemma will be a key tool to extend representations from algebra ideals.

Lemma 3.22. *Let $\mathcal{B}$ be a real involutive algebra and $\mathcal{A} \trianglelefteq \mathcal{B}$ be an ideal for which $(\mathcal{A}, p)$ is a seminormed involutive algebra. If $\pi\colon \mathcal{A} \to B(\mathcal{H})$ is a non-degenerate linear $*$-representation bounded on $\mathtt{ball}(\mathcal{A}, p)$ and $S := \{b \in \mathcal{B}\colon b\,\mathtt{ball}(\mathcal{A}, p) \cup \mathtt{ball}(\mathcal{A}, p)b \subseteq \mathtt{ball}(\mathcal{A}, p)\}$ generates $\mathcal{B}$ linearly, then there exists a unique linear representation $\widehat{\pi}\colon \mathcal{B} \to B(\mathcal{H})$ satisfying*

$$\widehat{\pi}(b)\pi(a) = \pi(ba) \quad \text{for} \quad a \in \mathcal{A}, b \in \mathcal{B}. \tag{5}$$

This representation is bounded on S.

Proof. In view of Lemma 3.13, the bounded representation $\pi|_{\mathtt{ball}(\mathcal{A},p)}$ extends to a bounded representation $\widetilde{\pi}\colon S \to B(\mathcal{H})$ which is uniquely determined by the relation

$$\widetilde{\pi}(s)\pi(a) = \pi(sa) \quad \text{for} \quad s \in S, a \in \mathtt{ball}(\mathcal{A}, p).$$

On the other hand, by [Ne00, Prop. II.4.14], there is a $*$-representation $\widehat{\pi}\colon \mathcal{B} \to \mathrm{End}(\mathcal{H}^0)$ on the dense subspace $\mathcal{H}^0 := \operatorname{span}\pi(\mathcal{A})\mathcal{H}$ of $\mathcal{H}$ which is uniquely determined by the relation $\widehat{\pi}(b)\pi(a)v = \pi(ba)v$ for $a \in \mathcal{A}, b \in \mathcal{B}, v \in \mathcal{H}$. The uniqueness and the linearity of π now imply that $\widehat{\pi}$ is linear. Since S spans $\mathcal{B}$ and $\widetilde{\pi}(s)|_{\mathcal{H}^0} = \widehat{\pi}(s)$ for $s \in S$, it further follows that all operators $\widehat{\pi}(b)$, $b \in \mathcal{B}$, are bounded, so that we actually obtain a $*$-representation $\widehat{\pi}\colon \mathcal{B} \to B(\mathcal{H})$ satisfying (5). □

We shall also need the following result.

Lemma 3.23. *Consider the abelian multiplicative semigroup $\mathcal{I} = ((0,1), \cdot)$ endowed with the trivial involution. Then the following assertions hold:*

(i) *If $\mathcal{B}$ is a C^*-algebra and $\varphi\colon \mathcal{I} \to \mathcal{B}$ is a positive definite function, then, for every $s \in \mathcal{I}$, we have $0 \leq \varphi(s) \in \mathcal{B}$.*

(ii) *If, moreover, $\mathcal{B} = \mathbb{C}$ and φ is bounded, then φ is also increasing and continuous.*

(iii) *Every bounded $*$-representation $(\pi, \mathcal{H})$ of $\mathcal{I}$ is norm-continuous and extends to a strongly continuous representation $\widehat{\pi}$ of the unital semigroup $(0, 1]$. Moreover, if π is non-degenerate and $V \subseteq \mathcal{H}$ is dense, then $\pi(r)V$ is dense for every $r \in \mathcal{I}$, and $\widehat{\pi}(1) = \mathbf{1}$.*

(iv) *Let V be a complex Hilbert space and $(A_n)_{n\geq 0}$ in $B(V)$ be a sequence for which the series $\varphi(t) = \sum_{n\geq 0} t^n A_n$ is norm convergent in $B(V)$ for every $t \in \mathcal{I}$ and the function $\varphi\colon \mathcal{I} \to B(V)$ is positive definite. Then $0 \leq A_n \leq \sup_{0<t\leq 1} \|\varphi(t)\| \cdot \mathbf{1}$ for every $n \geq 0$.*

Proof. (i) follows from $\varphi(s) = \varphi((s^{1/2})^* s^{1/2}) \geq 0$ for every $s \in \mathcal{I}$.

(ii) From (i) we know that $\varphi(\mathcal{I}) \subseteq [0,\infty)$. Now the conclusion follows by [Arv87, Prop. 2.4].

(iii) Since we may w.l.o.g. assume that π is non-degenerate and $\mathcal{I} \cong ((0,\infty),+)$, the first assertion follows from [Ne00, Lemma VI.2.2].

Assume that π is non-degenerate and that $V \subseteq \mathcal{H}$ is dense. If $w \in \mathcal{H}$ is orthogonal to $\pi(r)V$ for some $r \in \mathcal{I}$, then $\langle \pi(r)w, V\rangle = \langle w, \pi(r)V\rangle = \{0\}$, which leads to $\pi(r)w = 0$. In view of the Spectral Representation $\pi(r) = \int_0^\infty r^\alpha \, dP(\alpha)$ (see [RSzN72, Ch. X, §2, subsect. 141]), the fact that $r^\alpha > 0$ for every $\alpha \geq 0$ implies that $\pi(r)$ is injective, and therefore $w = 0$. This shows that $\pi(r)V$ is dense for every $r \in \mathcal{I}$. We also obtain $\widehat{\pi}(1) = \lim_{r\to 1} \pi(r) = \int_0^\infty dP(\alpha) = \mathbf{1}$.

(iv) For every vector $v \in V$, the function $\varphi^v\colon \mathcal{I} \to \mathbb{C}, s \mapsto \langle \varphi(s)v, v\rangle$ is positive definite and by (iii) it extends continuously to a positive definite function on $[0,1]$. Hence [Arv87, Lemma 3.8] shows that $0 \leq \langle A_n v, v\rangle \leq \sup_{0\leq t\leq 1} \langle \varphi(t)v, v\rangle$ for every $n \geq 0$, which implies the assertion. □

Finally, here is the **proof of Theorem 3.18:**

Proof. We shall prove the dilatability of φ by verifying that, for every $v \in V$, the function $\varphi \cdot v$ belongs to $\mathcal{H}_\varphi$ (Proposition 3.7). Proposition 3.21 applies in particular to structured $*$-semigroups because the subsemigroup $\mathcal{I} := \{\gamma(r)\colon 0 < r < 1\}$ of S° satisfies

$$S^\circ = S\mathcal{I} = \mathcal{I}S \quad \text{and} \quad S^\circ = S^\circ S^\circ$$

(cf. Remark 3.16). By construction, the space $\mathcal{H} := \mathcal{H}_\varphi \subseteq V^{S^\circ}$ consists of functions on S°. This subsemigroup satisfies $\mathcal{I}\mathcal{S}^\circ = \mathcal{S}^\circ = \mathcal{S}^\circ\mathcal{I}$, which implies that the representation $\pi_\varphi \circ \gamma$ of $(0,1)$ is also non-degenerate. From [Ne00, Lemma VI.2.2] it now follows that this representation of $((0,1),\cdot) \cong ((0,\infty),+)$ is norm-continuous and extends to a strongly continuous representation of the unital semigroup $(0,1]$.

For $v \in V$ and $s \in S^\circ$, we have

$$(K_s^* v)(t) = \varphi(ts^*)v \quad \text{and} \quad \pi_\varphi(s)K_t^* = K_{ts^*}^*. \tag{6}$$

For $v, w \in V$ and $s, t \in S^0$, we thus obtain

$$\langle \varphi(st^*)v, w\rangle = \langle K_t^* v, K_s^* w\rangle = \lim_{r\to 1-} \langle \pi_\varphi(r)K_t^* v, K_s^* w\rangle = \lim_{r\to 1-} \langle \varphi(srt^*)v, w\rangle.$$

Since $S^\circ = S^\circ\mathcal{I} = S^\circ S^\circ$, this implies that

$$\varphi(s) = \lim_{r\to 1-} \varphi(sr) \quad \text{for} \quad s \in S^\circ \tag{7}$$

in the weak operator topology. From Lemma 3.23 we obtain the relation

$$\varphi(\gamma(rr')) \leq \varphi(\gamma(r)) \quad \text{for} \quad 0 < r, r' < 1,$$

so that the boundedness of φ on $\mathcal{I}$ implies the existence of the strong operator limit

$$A := \lim_{r\to 1-} \varphi(\gamma(r)) \in B(V).$$

For $v \in V$ and $0 < r < s < 1$, we further get

$$\begin{aligned}\|K^*_{\gamma(r)}v - K^*_{\gamma(s)}v\|^2 &= \|K^*_{\gamma(r)}v\|^2 + \|K^*_{\gamma(s)}v\|^2 - 2\,\mathrm{Re}\langle K^*_{\gamma(r)}v, K^*_{\gamma(s)}v\rangle \\ &= \langle\varphi(\gamma(r^2))v, v\rangle + \langle\varphi(\gamma(s^2))v, v\rangle - 2\,\mathrm{Re}\langle\varphi(\gamma(rs))v, v\rangle \to 0\end{aligned}$$

for $r, s \to 1-$. This implies that the limit

$$\iota(v) := \lim_{r\to 1-} K^*_{\gamma(r)}v = \varphi \cdot v$$

exists for every $v \in V$ in $\mathcal{H}_\varphi$ with

$$\|\iota(v)\|^2 = \lim_{r\to 1-} \|K^*_{\gamma(r)}v\|^2 = \lim_{r\to 1-} \langle K(\gamma(r), \gamma(r))v, v\rangle = \lim_{r\to 1-} \langle\varphi(\gamma(r^2))v, v\rangle = \langle Av, v\rangle,$$

which leads to $\|\iota\|^2 \le \|A\| \le \limsup \|\varphi(\gamma(r))\|$. In particular, $\varphi \cdot v \in \mathcal{H}_\varphi$ holds for every $v \in V$, so that the assertion follows from Proposition 3.4. □

3.3 Extreme points and homogeneity

For later use we also state a rather standard result on the situation when the representation π_φ in Theorem 3.18 is irreducible (compare [Ne00, Cor. III.1.10] or [Arv87, Lemma 5.3]). We will use the following terminology.

Definition 3.24. If S is a structured $*$-semigroup and E is a real vector space, then a function $f\colon S^\circ \to E$ is said to be *homogeneous of degree* $\alpha \in \mathbb{R}$ if, for every $x \in S^\circ$ and $0 < r < 1$, we have $f(rx) = r^\alpha f(x)$.

Proposition 3.25. *Let S be a structured $*$-semigroup and $\mathrm{P}_1(S^\circ)$ the set of all positive definite functions $\varphi\colon S^\circ \to \mathbb{C}$ satisfying $\sup_{x\in S^\circ} |\varphi(x)| \le 1$. Then $\mathrm{P}_1(S^\circ)$, endowed with the topology of pointwise convergence, is a compact convex set. For every non-zero extreme point $\varphi \in \mathrm{P}_1(S^\circ)$, its corresponding representation $\pi_\varphi\colon S^\circ \to C(\mathcal{H}_\varphi)$ given by* Theorem 3.18 *is irreducible and there exists some $\alpha \in [0, \infty)$ for which φ is homogeneous of degree α and*

$$\pi_\varphi(\gamma(r)) = r^\alpha \mathbf{1} \quad \text{for} \quad 0 < r < 1.$$

If φ is non-constant and $t \mapsto \varphi(tx)$ is continuous on $[0, 1)$ for all $x \in S^\circ$, then $\alpha > 0$.

Proof. Let $\mathbb{D} := \{z \in \mathbb{C}\colon |z| \le 1\}$. Since positive definiteness is preserved under pointwise limits, $\mathrm{P}_1(S^\circ)$ is a closed subset of the compact set $\mathbb{D}^{S^\circ}$, hence compact.

Let $\varphi \in \mathrm{P}_1(S^\circ)$ be an extreme point. In view of Lemma 3.20, the representation $(\pi_\varphi, \mathcal{H}_\varphi)$ from Theorem 3.18 is non-degenerate and $v := \iota(1)$ satisfies

$$\|v\|^2 = \|\iota^*\iota\| \le \sup_{0<r<1} \|\varphi(r)\| \le 1.$$

Suppose that we have a direct sum decomposition $\pi_\varphi = \pi_1 \oplus \pi_2$ with $\mathcal{H}_\varphi = \mathcal{H}_1 \oplus \mathcal{H}_2$ and both summands non-zero. Then $v = (v_1, v_2) := \iota(1)$ is a cyclic vector with $\varphi(s) = \langle \pi_\varphi(s)v, v\rangle$ for $s \in S$. As $|\varphi(s)| \leq \|v\|^2 \leq 1$ and φ is a non-zero extreme point, we have $\|v\| = 1$. The cyclicity of v implies that both components v_j are non-zero. Accordingly

$$\varphi = \|v_1\|^2\varphi_1 + \|v_2\|^2\varphi_2 \quad \text{with} \quad \varphi_j(s) := \frac{\langle \pi_j(s)v_j, v_j\rangle}{\|v_j\|^2}$$

is a proper convex combination of the two elements $\varphi_j \in \mathrm{P}_1(S^\circ)$. This contradiction implies that π_φ is irreducible.

Now Schur's Lemma implies the existence of a function $c\colon (0,1) \to [0,1]$ with $\pi(\gamma(r)) = c(r)\mathbf{1}$ for $0 < r < 1$. From [Ne00, Lemma VI.2.2] it now follows that $\pi\circ\gamma$ is a norm-continuous representation, so that there exists some $\alpha \geq 0$ with $c(r) = r^\alpha$ for all $0 < r < 1$. We thus obtain

$$\varphi(rs) = \iota^*\pi(rs)\iota = \iota^*\pi(\gamma(r))\pi(s)\iota = r^\alpha\iota^*\pi(s)\iota = r^\alpha\varphi(s) \quad \text{for} \quad s \in S^\circ, 0 < r < 1.$$

This proves the first assertion, and the second follows directly. □

The following example shows that Proposition 3.25 does not generalize in any obvious way to operator-valued functions.

Example 3.26. Let $S := \{z \in \mathbb{C}\colon |z| \leq 1\} = C(\mathbb{C})$, $S^\circ = \{z \in \mathbb{C}\colon |z| < 1\}$ and, for a complex Hilbert space V, write $\mathrm{P}_1(S^\circ, B(V))$ the set of all positive definite functions $\varphi\colon S^\circ \to B(V)$ satisfying $\sup\limits_{x\in S^\circ} \|\varphi(x)\| \leq 1$. Then $\mathrm{P}_1(S^\circ, B(V))$, endowed with the topology of pointwise weak convergence, is a compact convex set.

The function

$$\varphi\colon S^\circ \to M_2(\mathbb{C}) \cong B(\mathbb{C}^2), \quad \varphi(z) = \begin{pmatrix} 1 & 0 \\ 0 & z \end{pmatrix}$$

is positive definite with $\|\varphi(s)\| = 1$ for every $s \in S^\circ$. The corresponding space $\mathcal{H}_\varphi \subseteq (\mathbb{C}^2)^{S^\circ}$ is generated by the function

$$K_{s,v}(t) = \varphi(ts^*)v = (v_1, ts^*v_2) \quad \text{for} \quad v = (v_1, v_2).$$

It follows that

$$\mathcal{H}_\varphi \cong \mathbb{C}\begin{pmatrix} 1 \\ 0 \end{pmatrix} \oplus \mathbb{C}z\begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

is 2-dimensional and multiplicity free.

If $\varphi = \lambda\varphi_1 + (1-\lambda)\varphi_2$ with $0 < \lambda < 1$ and $\varphi_1, \varphi_2 \in \mathrm{P}_1(S^\circ, B(\mathbb{C}^2))$, then the functions φ_j correspond to subrepresentations, so that the only non-trivial way to decompose φ is by

$$\lambda\varphi_1(z) = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \quad \text{and} \quad (1-\lambda)\varphi_2(z) = \begin{pmatrix} 0 & 0 \\ 0 & z \end{pmatrix}.$$

This leads to $\|\varphi_1(z)\| = \lambda^{-1}$, so that $\varphi_1 \notin \mathrm{P}_1(S^\circ, B(\mathbb{C}^2))$. Consequently, $\varphi \in \mathrm{P}_1(S^\circ, B(\mathbb{C}^2))$ is an extreme point, the corresponding representation is not irreducible and φ is not homogeneous.

4 Completely positive functions on $\mathtt{ball}(\mathcal{A}, p)$

In this section we first show that a dilation of a positive definite function on $\mathtt{ball}(\mathcal{A}, p)$ is completely positive if and only if φ is. Then we turn to the important special case $\mathcal{A} = \mathbb{R}$ which provides the key tools to obtain a series expansion $\varphi = \sum_{n=0}^{\infty} \varphi_n$ of a bounded completely positive function φ on $\mathtt{ball}(\mathcal{A}, p)$ into homogeneous components φ_n (Subsection 4.3).

4.1 Complete positivity of dilations

The following proposition connects the complete positivity of a function $\varphi\colon \mathtt{ball}(\mathcal{A}, p) \to B(V)$ to the complete positivity of the corresponding GNS representation $(\pi_\varphi, \mathcal{H}_\varphi)$. If $\mathcal{A}$ is unital, then φ is dilatable by Theorem 3.18, so that it asserts in this case that the complete positivity of φ and π_φ are equivalent. It generalizes [Arv87, Th. 2.17], which deals with the case $\dim V = 1$ and a unital C^*-algebra $\mathcal{A}$.

Proposition 4.1. *Let $(\mathcal{A}, p)$ be a seminormed $*$-algebra, V be a complex Hilbert space, and $\varphi\colon \mathtt{ball}(\mathcal{A}, p) \to B(V)$ be a bounded positive definite function. Then the following assertions hold:*

(a) *If φ is completely positive, then the corresponding GNS representation $(\pi_\varphi, \mathcal{H}_\varphi)$ is completely positive.*

(b) *If $(\pi, \mathcal{H}, \iota)$ is a minimal dilation of φ, then φ is completely positive if and only if π is completely positive.*

Proof. First note that for every $a \in \mathtt{ball}(\mathcal{A}, p)$ one has $\pi_\varphi(a) \in B(\mathcal{H}_\varphi)$ by Remark 3.5(a).

(a) Put $(\pi, \mathcal{H}) := (\pi_\varphi, \mathcal{H}_\varphi)$. We have to show that, if $n \geq 1$, $A := (a_{jk}) \in M_n(\mathcal{A})_+$ and $a_{jk} \in \mathtt{ball}(\mathcal{A}, p)$ for $1 \leq j, k \leq n$, then $(\pi(a_{jk})) \in M_n(B(\mathcal{H}))_+$. Since the functions $K_s^* v$, $s \in \mathtt{ball}(\mathcal{A}, p)$, $v \in V$, span a dense subspace of $\mathcal{H}$, it suffices to prove that $\sum_{j,k=1}^{n} \langle \pi(a_{jk}) h_k, h_j \rangle \geq 0$ for arbitrary $h_1, \ldots, h_n \in \operatorname{span}\{K_s^* v\colon s \in \mathtt{ball}(\mathcal{A}, p), v \in V\}$.

There exists an $m \in \mathbb{N}$, and $b_{jp} \in \mathtt{ball}(\mathcal{A}, p)$ and $v_{jp} \in V$ for $1 \leq j \leq n$ and $1 \leq p \leq m$, such that $h_j = \sum_{p=1}^{m} K_{b_{jp}}^* v_{jp}$. Then

$$\begin{aligned}\sum_{j,k=1}^{n} \langle \pi(a_{jk}) h_k, h_j \rangle &= \sum_{j,k=1}^{n} \sum_{p,q=1}^{m} \langle \pi(a_{jk}) K_{b_{kp}}^* v_{kp}, K_{b_{jq}}^* v_{jq} \rangle \\ &= \sum_{j,k=1}^{n} \sum_{p,q=1}^{m} \langle K_{b_{jq}} \pi(a_{jk}) K_{b_{kp}}^* v_{kp}, v_{jq} \rangle = \sum_{j,k=1}^{n} \sum_{p,q=1}^{m} \langle \varphi(b_{jq} a_{jk} b_{kp}^*) v_{kp}, v_{jq} \rangle,\end{aligned}$$

and since φ is completely positive, it suffices to check that the matrix

$$C := (b_{jq} a_{jk} b_{kp}^*) \in M_{mn}(\mathcal{A})$$

is non-negative. If we denote by $B_p \in M_n(\mathcal{A})$ the diagonal matrix $\operatorname{diag}(b_{1p}, \dots, b_{np})$, then $C = D(A \otimes \mathbf{1}_m)D^* \geq 0$, where

$$D = \begin{pmatrix} B_1 & 0 & \dots & 0 \\ B_2 & 0 & \dots & 0 \\ \vdots & & \vdots & \\ B_m & 0 & \dots & 0 \end{pmatrix} \in M_m(M_n(\mathcal{A})) \cong M_{mn}(\mathcal{A})$$

and this proves that π is completely positive.

(b) It is well known (and easy to see) that the linear mapping $\Psi\colon B(\mathcal{H}) \to B(V), T \mapsto \iota^* T \iota$ is completely positive. Therefore $\varphi = \Psi \circ \pi$ is completely positive if π is completely positive. For the converse we can use (a) because $(\pi, \mathcal{H})$ is equivalent to the GNS representation $(\pi_\varphi, \mathcal{H}_\varphi)$ by Proposition 3.4. □

Corollary 4.2. *A bounded $*$-representation $(\pi, \mathcal{H})$ of $\mathtt{ball}(\mathcal{A}, p)$ is completely positive if and only if, for every v in some dense subset of $\mathcal{H}$, the function $\pi^v(a) := \langle \pi(a)v, v\rangle$ is completely positive.*

Proof. If π is completely positive, then so is every function π^v since the linear functional $A \mapsto \langle Av, v\rangle$ on $B(\mathcal{H})$ is positive definite, hence completely positive (Remark 2.10(c)).

Suppose, conversely, that all functions π^v are completely positive for v in some dense subset of $\mathcal{H}$, hence actually for every $v \in \mathcal{H}$. Write π as the direct sum of a zero representation and a direct sum of cyclic representations $(\pi_j, \mathcal{H}_j, v_j)_{j\in J}$. Since $v_j \in \mathcal{H}_j$ is a cyclic vector for the representation π_j, it follows that π_j is a minimal dilation of the positive definite function $\varphi_j := \pi^{v_j}$, and then π_j is unitarily equivalent to the GNS representation π_{φ_j} by Proposition 3.4(b). On the other hand, since the function φ_j is completely positive by assumption, the representation π_{φ_j} is completely positive by Proposition 4.1(a). Hence also π_j is completely positive. This implies that π is completely positive, as an orthogonal direct sum of completely positive maps. □

Corollary 4.3. *Let $(\mathcal{A}, p)$ be a seminormed $*$-algebra and V be a complex Hilbert space. If $\varphi\colon \mathtt{ball}(\mathcal{A}, p) \to B(V)$ is a dilatable completely positive function, then $\varphi - \varphi(0)$ is also completely positive and dilatable.*

Proof. Let $(\pi, \mathcal{H}, \iota)$ be a dilation of φ. With the same argument as in the proof of Lemma 3.6, we see that $P := \pi_\varphi(0)$ is a projection onto a trivial subrepresentation. With $\mathcal{H}_0 := P(\mathcal{H})$ and $\mathcal{H}_1 := \mathcal{H}_0^\perp$ and the corresponding decompositions $\iota = \iota_0 + \iota_1$ and $\pi(s) = (\pi_0(s), \pi_1(s)) = (\mathbf{1}, \pi_1(s))$, we then obtain

$$\varphi(s) = \iota^*\pi(s)\iota = \iota_0^*\iota_0 + \iota_1^*\pi_1(s)\iota_1 \quad \text{and} \quad \varphi(0) = \iota^*\pi(0)\iota = \iota_0^*\iota_0.$$

This proves the assertion. □

Lemma 4.4. *Let $(\pi, \mathcal{H})$ be a non-degenerate bounded representation of $\mathtt{ball}(\mathcal{A}, p)$, with its canonical extension $\pi^1\colon \mathtt{ball}(\mathcal{A}^1, p^1) \to B(\mathcal{H})$ (Example 3.14). Then π is completely positive if and only if π^1 is completely positive.*

Proof. We must show that the complete positivity of π implies the complete positivity of π^1. Decomposing π as a direct sum of cyclic representations, it suffices to prove the lemma for cyclic representation. So assume that $0 \neq v \in \mathcal{H}$ is cyclic and put $\varphi(a) := \langle \pi(a)v, v\rangle$. For $s_1, \ldots, s_n \in \mathtt{ball}(\mathcal{A}, p)$ and $c_1, \ldots, c_n \in \mathbb{C}$, the vectors of the form $w := \sum_{j=1}^n c_j \pi(s_j)v$ form a dense subspace of $\mathcal{H}$. In view of Corollary 4.2, it therefore suffices to show that the functions

$$\psi\colon \mathtt{ball}(\mathcal{A}^1, p^1) \to \mathbb{C}, \quad \psi(a) := \langle \pi^1(a)w, w\rangle = \sum_{i,j=1}^n c_i\overline{c_j}\langle \pi(s_j^* a s_i)v, v\rangle = \sum_{i,j=1}^n c_i\overline{c_j}\varphi(s_j^* a s_i)$$

are completely positive.

So let $A = (a_{k\ell}) \in M_N(\mathcal{A}^1)$ be positive. Then the matrix $B := (s_j^* a_{k\ell} s_i) \in M_{nN}(\mathcal{A})$ is positive by Example 2.17(a). Now the complete positivity of φ implies that the matrix $(\varphi(s_j^* a_{k\ell} s_j)) \in M_{nN}(\mathbb{C})$ is positive, so that Example 2.17(b) shows that the matrix

$$\psi_N(A) = \Big(\sum_{i,j=1}^n c_i\overline{c_j}\varphi(s_j^* a_{k\ell} s_i)\Big)_{1\leq k,\ell\leq N} \in M_N(\mathbb{C})$$

is positive. We conclude that ψ is completely positive. □

The following proposition is a version of Proposition 3.25 for completely positive functions on $\mathtt{ball}(\mathcal{A}, p)$.

Proposition 4.5. *If $(\mathcal{A}, p)$ is a seminormed involutive algebra, then the set $\mathrm{CP}_1(\mathtt{ball}(\mathcal{A}, p))$ of all completely positive functions $\varphi\colon \mathtt{ball}(\mathcal{A}, p) \to \mathbb{C}$ with*

$$\|\varphi\|_\infty := \sup_{x\in \mathtt{ball}(\mathcal{A},p)} |\varphi(x)| \leq 1,$$

endowed with the topology of pointwise weak convergence, is a compact convex set. For every extreme point $\varphi \in \mathrm{CP}_1(\mathtt{ball}(\mathcal{A}, p))$ either $\mathcal{H}_\varphi = \{0\}$ or the representation $\pi_\varphi\colon S^\circ \to C(\mathcal{H}_\varphi)$ is irreducible and there exists some integer $\alpha \in \mathbb{N}_0$ such that

$$\varphi(tab) = t^\alpha \varphi(ab) \quad \text{for} \quad 0 \leq |t| < 1, a, b \in \mathtt{ball}(\mathcal{A}, p). \tag{8}$$

Proof. Note that complete positivity is preserved under pointwise convergence, so that the same argument as in Proposition 3.25 implies the compactness of $\mathrm{CP}_1(\mathtt{ball}(\mathcal{A}, p))$. We likewise see that, for every extreme point φ of $\mathrm{CP}_1(\mathtt{ball}(\mathcal{A}, p))$, for which the corresponding Hilbert space $\mathcal{H}_\varphi$ is non-zero, the corresponding representation π_φ is irreducible. Here we use that the complete positivity of φ implies the complete positivity of π_φ (Proposition 4.1(a)), and therefore the complete positivity of the functions $\varphi_j(s) := \frac{\langle \pi_j(s)v_j, v_j\rangle}{\|v_j\|^2}$. Using the canonical extension of the representation to $\mathtt{ball}(\mathcal{A}^1, p^1)$ (Example 3.14) and Schur's Lemma, this implies the existence of some $\alpha \geq 0$ with $\widetilde{\pi}_\varphi(t) = t^\alpha \mathbf{1}$ for $|t| < 1$. For $v \in V$ and $a \in \mathtt{ball}(\mathcal{A}, p)$, we now have

$$t^\alpha K_{a,v}(b) = (\widetilde{\pi}_\varphi(t)K_{a,v})(b) = K_{a,v}(tb) = \varphi(tba^*)v,$$

so that φ satisfies (8). It remains to show that $\alpha \in \mathbb{N}_0$.

If $\langle\varphi(a^*a)v,v\rangle = 0$ for every $a \in \mathtt{ball}(\mathcal{A},p)$ and $v \in V$, then the relation $\|K_{a,v}\|^2 = \langle\varphi(a^*a)v,v\rangle$ in $\mathcal{H}_\varphi$ implies that $\mathcal{H}_\varphi = \{0\}$. If this is not the case, then there exists an $a \in \mathtt{ball}(\mathcal{A},p)$ and $v \in V$ such that the completely positive function $\psi(t) := \langle\varphi(a^*ta)v,v\rangle = t^\alpha\langle\varphi(a^*a)v,v\rangle$ on $[-1,1]$ does not vanish. Its complete positivity implies that, for every positive definite matrix $A = (a_{ij}) \in M_n(\mathbb{R})$ with $a_{ij} > 0$, the matrix $(a_{ij}^\alpha) \in M_n(\mathbb{R})$ is positive semidefinite. In view of [Ho69, pp. 270-271], this implies that $\alpha \in \mathbb{N}_0$ (see also [CR78, Thm. 1]). □

Proposition 4.6. *Let $(\mathcal{A},p)$ be a seminormed involutive algebra and $\varphi\colon \mathcal{A} \to B(V)$ be dilatable completely positive and homogeneous of degree $n \in \mathbb{N}_0$ and bounded on $\mathtt{ball}(\mathcal{A},p)$. Then φ is a continuous polynomial which is homogeneous of degree n.*

Proof. **Case 1:** First we assume that $\mathcal{A}$ is unital. We use the method of proof of [Arv87, Lemma 4.3] to see that, for $h \in \mathcal{A}$, the function

$$(\Delta_h\varphi)(a) := \varphi(a+h) - \varphi(a)$$

is also completely positive. Iterating this argument, one shows that the function

$$(\Delta^n\varphi)(a;h_1,\dots,h_n) := (\Delta_{h_1}\cdots\Delta_{h_n})(a)$$

is completely positive on the seminormed involutive algebra $\mathcal{A}^{n+1}$ with respect to the seminorm

$$q_{n+1}(a_0,\dots,a_n) := \max\{p(a_0),\cdots,p(a_n)\}.$$

The corresponding argument in the proof of [Arv87, Lemma 5.8] works for general unital involutive algebras, but one should replace the reference to [Arv87, Thm. 2.17] in its proof by a reference to [Arv87, Lemma 4.3].

Theorem 3.18 shows that, if $\varphi(r) = 0$ for $0 < r < 1$, then $\iota = 0$, and thus $\varphi = 0$. Now the same argument as in [Arv87, p. 342] implies that φ is a homogeneous polynomial of degree n. In the whole argument, continuity is never needed, only that the functions $\Delta^k\varphi$ are bounded on balls with respect to the corresponding seminorm q_{k+1}.

Case 2: Now we assume that $\mathcal{A}$ is not unital. Let $(\pi,\mathcal{H},\iota)$ be a dilation of φ. Since we may assume that $\mathcal{H} = \mathcal{H}_\varphi$ (Proposition 3.4) and φ is homogeneous of degree n, it follows that $\pi(ta) = t^n\pi(a)$ for $a \in \mathcal{A}$ and $t \in \mathbb{R}$. In view of Lemma 4.4, the bounded representation π of $\mathtt{ball}(\mathcal{A},p)$ extends to a bounded representation π^1 of $\mathtt{ball}(\mathcal{A}^1,p^1)$. Further Proposition 4.1 implies that π^1 is completely positive. Therefore the first part of the proof applies to the completely positive function $\varphi^1(a) := \iota^*\pi^1(a)\iota$ which extends φ to $\mathtt{ball}(\mathcal{A}^1,p^1)$. □

Since representations are dilatable for trivial reasons (Example 3.8), the preceding proposition implies:

Corollary 4.7. *If $\pi\colon (\mathcal{A},\cdot) \to B(\mathcal{H})$ is a representation bounded on $\mathtt{ball}(\mathcal{A},p)$ and homogeneous of degree n, then π is a continuous homogeneous polynomial of degree n.*

Remark 4.8. Let $\varphi\colon \mathcal{A} \to B(V)$ be completely positive, homogeneous of degree $n \in \mathbb{N}_0$ and bounded on $\mathtt{ball}(\mathcal{A},p)$ but not necessarily dilatable. Then all functions in $\mathcal{H}_\varphi \subseteq \mathbb{C}^{\mathtt{ball}(\mathcal{A},p)}$ are homogeneous of degree $n \in \mathbb{N}_0$, which implies that the representation π_φ is homogeneous of degree n. Now the preceding corollary implies that π_φ is a continuous homogeneous polynomial of degree n.

Remark 4.9. It follows by [AC86, Th. 4] that if $\mathcal{A}$ is a C^*-algebra and $S = (\mathcal{A}, \cdot)$ its multiplicative group, then the completely positive maps $\varphi\colon S \to B(V)$ which are homogeneous of degree 1 in the sense of the above Definition 3.24 are precisely the (completely positive) $\mathbb{R}$-linear maps.

4.2 The case $\mathcal{A} = \mathbb{R}$

In this section we take a closer look at the special case $\mathcal{A} = \mathbb{R}$. We write $S := \mathtt{ball}(\mathbb{R}) = ((-1,1), \cdot)$. We know from Proposition 4.5 that the set $\mathrm{CP}_1(S)$ of completely positive functions φ on S with $\|\varphi\|_\infty \leq 1$ is a compact convex set with respect to the topology of pointwise convergence and that its non-zero extreme points are the characters

$$\chi_n(r) = r^n, \quad n \in \mathbb{N}_0.$$

Then

$$\mathbb{S} := \Big\{(a_n)_{n\in\mathbb{N}_0} \in \ell^1(\mathbb{N}_0)\colon a_j \geq 0, \sum_n a_n \leq 1\Big\}$$

is a weak-$*$-closed bounded subset of $\ell^1(\mathbb{N}_0) \cong c_0(\mathbb{N}_0)'$, hence compact in the weak-$*$-topology. This implies that on $\mathbb{S}$ the weak-$*$-topology coincides with the product topology inherited from $\mathbb{R}^{\mathbb{N}_0}$. We have a well-defined map

$$\Phi\colon \mathbb{S} \to \mathrm{CP}_1((-1,1)), \quad \Phi((a_n))(t) := \sum_{n=0}^{\infty} a_n t^n,$$

and since the sequences $(t^n)_{n\in\mathbb{N}_0}$, $|t| < 1$, are in $c_0(\mathbb{N}_0)$, Φ is continuous with respect to the product topology on $\mathbb{S}$ and the topology of pointwise convergence on $\mathrm{CP}_1((-1,1))$. Since Φ is affine, its image is a compact convex subset of $\mathrm{CP}_1((-1,1))$ which contains the extreme points $\chi_n = \Phi(e_n)$. Hence Φ is surjective by the Krein–Milman Theorem. As Φ is obviously injective, the compactness of $\mathbb{S}$ implies that Φ is a homeomorphism. Hence every function $\varphi \in \mathrm{CP}_1((-1,1))$ is analytic and its Taylor series in 0 converges uniformly on the closed interval $[-1,1]$. In particular, φ extends to a continuous function on $[-1,1]$.

Remark 4.10. The completely positive characters of $(-1,1)$ are the monomials $\chi_n(t) = t^n$, $n \in \mathbb{N}_0$.

Proposition 4.11. (Completely positive functions on $(-1,1)$).

(a) *Every non-degenerate bounded completely positive representation $(\pi, \mathcal{H})$ of $(-1,1)$ is a direct sum of subrepresentations $(\pi_n, \mathcal{H}_n)_{n\in\mathbb{N}_0}$ with $\pi_n(t) = t^n\mathbf{1}$ for $|t| < 1$.*

(b) *Every bounded completely positive function $\varphi\colon (-1,1) \to B(V)$ is of the form*

$$\varphi(t) = \sum_{n=0}^{\infty} t^n A_n,$$

where $A_n \geq 0$ and the sum converges on each interval $[-r,r]$, $r < 1$, in the norm topology. Further, $\varphi(\pm 1) := \sum_{n=0}^{\infty} (\pm 1)^n A_n$ defines a weakly continuous extension to $[-1,1]$.

Proof. (a) Since every non-degenerate representation is a direct sum of cyclic ones, this follows from the structure of the bounded completely positive functions on $(-1,1)$:

$$\varphi(t) = \sum_{n=0}^{\infty} a_n t^n \quad \text{with} \quad a_n \geq 0 \quad \text{and} \quad \sum_n a_n < \infty.$$

It implies that $\mathcal{H}_\varphi$ contains χ_n whenever $a_n > 0$ because $\varphi - a_n\chi_n$ is positive definite ([Ne00, Thm. I.2.8]). Let $\varphi_n := a_n\chi_n$. Then the norm of φ_n in $\mathcal{H}_{\varphi_n}$ is given by $\|\varphi_n\|^2 = a_n$, so that the series $\sum_n \varphi_n$ converges in $\oplus_{n=0}^\infty \mathcal{H}_{\varphi_n}$. This shows that $\mathcal{H}_\varphi \cong \oplus_{n=0}^\infty \mathcal{H}_{\varphi_n}$.

(b) Let $(\pi, \mathcal{H}, \iota)$ be a dilation of φ (Theorem 3.18). Then Proposition 4.1 implies that the non-degenerate representation π is completely positive. Now (a) implies that $(\pi, \mathcal{H}) = \oplus_{n=0}^\infty (\pi_n, \mathcal{H}_n)$ and we obtain with $\iota(v) = \sum_n \iota_n(v)$, $\iota_n(v) \in \mathcal{H}_n$, the relations

$$\pi(t) = \sum_n t^n \operatorname{id}_{\mathcal{H}_n} \quad \text{and} \quad \varphi(t) = \sum_n t^n A_n \quad \text{with} \quad A_n := \iota_n^*\iota_n \in B(V),$$

where the series converge in the weak operator topology and uniformly on $[-r, r]$ in the norm topology. We further have the weak operator convergent series

$$\iota^*\iota = \sum_n \iota_n^*\iota_n = \sum_n A_n$$

which leads to weak operator continuous extension of φ by $\varphi(\pm 1) := \sum_{n=0}^\infty (\pm 1)^n A_n$. □

4.3 Series expansion of bounded completely positive functions

For a bounded function $\varphi\colon \mathtt{ball}(\mathcal{A}, p) \to E$ with values in a Banach space E, we write

$$\|\varphi\|_\infty := \sup\{\|\varphi(a)\| \colon a \in \mathtt{ball}(\mathcal{A}, p)\}.$$

Theorem 4.12. *Let $\varphi\colon \mathtt{ball}(\mathcal{A}, p) \to B(V)$ be a bounded dilatable completely positive function. Then φ is analytic in the p-topology and its Taylor series $\varphi = \sum_n \varphi_n$ at 0 converges for every $r < 1$ uniformly on the ball $\{a \in \mathcal{A} \colon p(a) \leq r\}$. The homogeneous polynomials φ_n are also completely positive with*

$$\|\varphi_n\|_\infty \leq \|\varphi\|_\infty \quad \textit{for every} \quad n \in \mathbb{N}.$$

Proof. Let $(\pi, \mathcal{H}, \iota)$ be a be a minimal dilation of φ (Theorem 3.18). Then the representation π is non-degenerate and is completely positive by Proposition 4.1, and by Lemma 4.4 it extends to a completely positive representation π^1 of $\mathtt{ball}(\mathcal{A}^1, p^1)$. Applying Proposition 4.11 to $\pi^1|_{(-1,1)\mathbf{1}}$, we see that $(\pi, \mathcal{H}) \cong \bigoplus_{n\in\mathbb{N}_0} (\pi_n, \mathcal{H}_n)$ with the $\mathtt{ball}(\mathcal{A}, p)$-invariant subspaces

$$\mathcal{H}_n = \{v \in \mathcal{H} \colon \pi^1(t\mathbf{1})v = t^n v \quad \text{for} \quad |t| < 1\}.$$

Writing $\iota(v) = \sum_n \iota_n(v)$ with $\iota_n(v) \in \mathcal{H}_n$, this leads to the weakly convergent series

$$\varphi(a) = \iota^*\pi(a)\iota = \sum_n \varphi_n(a) \quad \text{with} \quad \varphi_n(a) := \iota_n^*\pi_n(a)\iota_n,$$

where π_n and φ_n are homogeneous of degree n in the sense that

$$\varphi_n(ta) = t^n \varphi_n(a) \quad \text{and} \quad \pi_n(ta) = t^n \pi_n(a) \quad \text{for} \quad a \in \mathtt{ball}(\mathcal{A}, p), |t| < 1.$$

Since $\|\iota_n\| \leq \|\iota\|$ and $\|\pi_n(a)\| \leq \|\pi(a)\| \leq 1$, the series $\varphi(a) = \sum_{n=0}^{\infty} \varphi_n(a)$ converges uniformly on all subsets $B_r := \{a \in \mathcal{A} \colon p(a) < r\}$ for $r < 1$ with respect to the norm topology on $B(V)$, and pointwise on all of $\mathtt{ball}(\mathcal{A}, p)$. To see that φ is analytic, it now suffices to observe that each φ_n is polynomial, which follows from Lemmas 2.11 and Proposition 4.6. □

The main point of the preceding theorem is the fact that the homogeneous functions φ_n are polynomials. This would follow much more directly if φ is weakly smooth, because then the Taylor series of the functions $a \mapsto \langle \varphi(a)v, w\rangle$, $v, w \in V$, makes sense, and the terms of order n are polynomial in a.

Remark 4.13. (On the non-unital case) Every completely positive representation $\pi \colon \mathtt{ball}(\mathcal{A}, p) \to C(\mathcal{H})$ is in particular a dilatable completely positive function, so that Theorem 4.12 applies.

This applies in particular to the GNS representation $(\pi, \mathcal{H}) = (\pi_\varphi, \mathcal{H}_\varphi)$ associated to a bounded completely positive function $\varphi \colon \mathtt{ball}(\mathcal{A}, p) \to B(V)$ (Proposition 4.1 and Remark 3.5). From the decomposition $\pi = \sum_{n=0}^{\infty} \pi_n$, we derive that the corresponding positive definite kernel $K(a, b) := \varphi(ab^*)$ decomposes as $K = \sum_{n=0}^{\infty} K^n$, and these kernels satisfy

$$K^n(ab, c) = K^n(a, cb^*) \quad \text{for} \quad a, b, c \in \mathtt{ball}(\mathcal{A}, p).$$

If $\mathcal{A}$ is not unital, then it is not clear if there exists functions $\varphi_n \colon \mathtt{ball}(\mathcal{A}, p) \to B(V)$, $n \in \mathbb{N}_0$, with $\varphi_n(ab^*) = K^n(a, b)$ for $a, b \in \mathtt{ball}(\mathcal{A}, p)$. Without the assumption of dilatability it is therefore not clear if φ admits a corresponding series expansion. If $\mathtt{ball}(\mathcal{A}, p)$ contains an approximate identity and φ is continuous, then we can use [Ne00, Prop. IV.1.29] to obtain functions φ_n on the dense subsemigroup $\mathtt{ball}(\mathcal{A}, p)\,\mathtt{ball}(\mathcal{A}, p)$ of $\mathtt{ball}(\mathcal{A}, p)$ which satisfy $\varphi_n(ab^*) = K_n(a, b)$ for $a, b \in \mathtt{ball}(\mathcal{A}, p)$.

5 Analytic positive definite functions

In this section we turn to positive definite functions $\varphi \colon \mathtt{ball}(\mathcal{A}, p) \to B(V)$ which are analytic with respect to some locally convex topology on $\mathcal{A}$ for which p is a continuous seminorm (cf. Definition A.1). After some general observations concerning series expansions on structured semigroup in Subsection 5.1, we derive in Subsection 5.2 a series expansion of dilatable bounded analytic positive definite functions $\varphi \colon \mathtt{ball}(\mathcal{A}, p) \to B(V)$. Before we can use these result to show that φ is completely positive, we take in Subsection 5.3 a detailed look at the case where φ is a homogeneous polynomial. The main result to deal with this case is Proposition 5.11 which asserts that the C^*-algebra $S^n(C^*(\mathcal{A}, p))$ is the universal enveloping algebra of $(S^n(\mathcal{A}), p_n)$. This in turn is used to show that every dilatable positive definite homogeneous polynomial $\varphi \colon \mathcal{A} \to B(V)$ which is bounded on $\mathtt{ball}(\mathcal{A}, p)$ is completely positive.

5.1 Series expansions on structured $*$-semigroups

Proposition 5.1. *Let S be a structured $*$-semigroup, $T \subseteq S$ be a subsemigroup with $rT \subseteq T$ for $0 < r < 1$ and V be a complex Hilbert space. For every integer $n \geq 0$, let the function $\varphi_n \colon T \to B(V)$ be homogeneous of degree n such that, for every $a \in T$, the series*

$$\varphi(a) := \sum_{n \geq 0} \varphi_n(a)$$

is convergent in the weak operator topology and $M := \sup_{a \in T} \|\varphi(aa^*)\| < \infty$. Then the function φ is positive definite if and only if all the functions $(\varphi_n)_{n \geq 0}$ are positive definite on T. If this is the case, then $\sup_{a \in T} \|\varphi_n(aa^*)\| \leq M$ for all $n \geq 0$.

Proof. Let us fix arbitrary $m \geq 1$, $a_1, \ldots, a_m \in T$, and $v_1, \ldots, v_m \in V$, and set $\alpha_n := \sum_{j,k=1}^{m} \langle \varphi_n(a_j^* a_k) v_k, v_j \rangle \in \mathbb{C}$. If $0 \leq t \leq 1$, the series $\psi(t) = \sum_{n \geq 0} \alpha_n t^n = \sum_{j,k=1}^{m} \langle \varphi(t a_j^* a_k) v_k, v_j \rangle$ is convergent. Moreover, if $0 \leq t_1, \ldots, t_q \leq 1$, and $\mu_1, \ldots, \mu_q \in \mathbb{C}$, then

$$\sum_{i,\ell=1}^{q} \psi(t_i t_\ell) \mu_i \overline{\mu}_\ell = \sum_{i,\ell=1}^{q} \sum_{j,k=1}^{m} \langle \varphi((t_\ell a_j)^* (t_i a_k))(\mu_i v_k), \mu_\ell v_j \rangle \geq 0$$

since φ is positive definite. Therefore $\psi \colon [0,1] \to \mathbb{C}$ is positive definite on the $*$-semigroup $[0,1]$, and then [Arv87, Lemma 3.8] shows that, for every $n \geq 0$, we have

$$0 \leq \alpha_n \leq \sup_{0 \leq t \leq 1} \psi(t),$$

that is

$$0 \leq \sum_{j,k=1}^{m} \langle \varphi_n(a_j^* a_k) v_k, v_j \rangle \leq \sup_{0 \leq t \leq 1} \sum_{j,k=1}^{m} \langle \varphi(t a_j^* a_k) v_k, v_j \rangle. \tag{9}$$

Recalling the definition of α_n, it follows that, for each $n \geq 0$, the homogeneous function $\varphi_n \colon T \to B(V)$ is positive definite. Using the above equation (9) (with $m = 1$), we obtain

$$\sup_{a \in T} \|\varphi_n(a^* a)\| \leq \sup_{a \in T} \|\varphi(a^* a)\| = M.$$

This concludes the proof. □

Corollary 5.2. *Let S be a structured $*$-semigroup and V be any complex Hilbert space. For every integer $n \geq 0$, let the function $\varphi_n \colon S^\circ \to B(V)$ be homogeneous of degree n such that, for every $a \in S^\circ$, the series*

$$\varphi(a) := \sum_{n \geq 0} \varphi_n(a)$$

is convergent in the weak operator topology and $M := \sup_{a \in S^\circ} \|\varphi(a)\| < \infty$. Then the function φ is positive definite if and only if all the functions $(\varphi_n)_{n \geq 0}$ are positive definite. If this is the case, then $\sup_{a \in S^\circ} \|\varphi_n(a)\| \leq M$ for all $n \geq 0$.

Proof. We apply Proposition 5.1 to $T = S^\circ$ and recall that $S^\circ = S^\circ S^\circ$, which implies that, for $s = ab^*$, we obtain

$$\|\varphi_n(s)\|^2 = \|\varphi_n(ab^*)\|^2 \leq \|\varphi_n(aa^*)\|\|\varphi_n(bb^*)\| \leq M^2.$$

This completes the proof. □

5.2 Analytic positive definite functions

Definition 5.3. If M is an analytic manifold and V a complex Hilbert space, then we call a function $\varphi\colon M \to B(V)$ *weakly analytic* if there exists a dense subspace $E \subseteq V$, such that, the function $M \to \mathbb{C}, m \mapsto \langle \varphi(m)v, w\rangle$ is analytic for $v, w \in E$.

Proposition 5.4. *Every bounded weakly analytic positive definite function $\varphi\colon (-1,1) \to B(V)$ is completely positive.*

Proof. (a) First we consider the case $V = \mathbb{C}$. Then $\varphi|_{[0,1)}$ is increasing by Lemma 3.23(ii), and by [Ne00, Cor. VI.2.11] there exists a uniquely determined positive Radon measure μ on $[0,\infty)$ with

$$\varphi(t) = \int_0^\infty t^\alpha \, d\mu(\alpha) \quad \text{for} \quad 0 < t < 1.$$

On the other hand, the analyticity of φ in 0 implies the existence of an $\varepsilon > 0$ and $a_n \in \mathbb{C}$ with

$$\varphi(t) = \sum_{n=0}^\infty a_n t^n \quad \text{for} \quad |t| \leq \varepsilon.$$

Then the function $\psi(t) := \varphi(t\varepsilon) = \sum_{n=0}^\infty a_n \varepsilon^n t^n$ on $[0,1]$ is is positive definite, so that Lemma 3.23(iv) implies that $0 \leq a_n \leq \varepsilon^{-n}\psi(1) = \varepsilon^{-n}\varphi(\varepsilon)$. From the uniqueness of the measure μ and the identity

$$\int_0^\infty t^\alpha \, \varepsilon^\alpha d\mu(\alpha) = \varphi(t\varepsilon) = \sum_{n=0}^\infty a_n \varepsilon^n t^n \quad \text{for} \quad 0 < t \leq 1,$$

it follows that $\mu = \sum_{n=0}^\infty a_n \delta_n$. This in turn implies that

$$\varphi(t) = \int_0^\infty t^\alpha \, d\mu(\alpha) = \sum_{n=0}^\infty a_n t^n$$

holds for $0 \leq t < 1$ and by analyticity we have this relation for $|t| < 1$. It follows in particular that φ is completely positive.

(b) Now let $\varphi\colon (-1,1) \to B(V)$ be weakly analytic and positive definite. Let $E \subseteq V$ be a dense subspace such that $s \mapsto \langle \varphi(s)v, w\rangle$ is analytic for $v, w \in E$. To see that φ is completely positive, it suffices to show that the corresponding GNS representation π_φ is completely positive (Proposition 4.1). This representation is weakly analytic, because the functions

$$s \mapsto \langle \pi_\varphi(s)\pi_\varphi(t)\iota(v), \pi_\varphi(u)\iota(w)\rangle = \langle \varphi(ust)v, w\rangle, \qquad v, w \in E,$$

are analytic for every $t, u \in (-1,1)$. Therefore we may assume that $\varphi = \pi$ is a non-degenerate representation. Then π is a direct sum of cyclic representations whose generating vectors have analytic matrix coefficients. Combining (a) with Proposition 4.1, it follows that all these subrepresentations are completely positive, and therefore π is completely positive. □

Let $\varphi\colon S := \mathtt{ball}(\mathcal{A}, p) \to B(V)$ be a bounded positive definite real analytic function for some locally convex topology for which p is continuous (Definition A.1(c)) and $\pi_\varphi\colon S \to B(\mathcal{H}_\varphi)$ be the corresponding contraction representation of S (Theorem 3.18). Since φ is analytic, there exists an open circular 0-neighborhood $U \subseteq S$ such that

$$\varphi(a) = \sum_{n\geq 0} \varphi_n(a) \quad \text{with} \quad \varphi_n(a,\dots,a) = \frac{1}{n!}(\partial_a^n \varphi)(0)$$

converges for every $a \in U$.

Lemma 5.5. *If the function φ is dilatable, then the Hilbert space $\mathcal{H}_\varphi$ is the orthogonal direct sum of the subspaces $\mathcal{H}_{\varphi_n}$ and the series $\varphi = \sum_n \varphi_n$ converges uniformly on every ball $B_r := \{a \in \mathcal{A}\colon p(a) < r\}$.*

Proof. Let $\mathcal{H} := \mathcal{H}_\varphi$ and write π for the corresponding representation of S on $\mathcal{H}$ defined by $(\pi(s)f)(t) = f(ts)$. We write π^1 for the canonical extension of π to $\mathtt{ball}(\mathcal{A}^1, p^1)$ (Example 3.14). For $w := K_{s,v}$ and $t \in \mathcal{J} := (-1,1)$, we have

$$\langle \pi^1(t)w, w\rangle = \langle \pi^1(t)K_{s,v}, K_{s,v}\rangle = \langle K_{s,v}(st), v\rangle = \langle K_{st}K_s^* v, v\rangle = \langle \varphi(sts^*)v, v\rangle,$$

and this function on $\mathcal{J}$ is analytic and positive definite, hence completely positive by Proposition 5.4. We conclude that w has in $\mathcal{H}$ an orthogonal expansion $w = \sum_n w_n$ with $\pi^1(t)w_n = t^n w_n$ for $t \in \mathcal{J}$. As the vectors of the form $K_{s,v}$, $s \in S$, $v \in V$, span a dense subspace of $\mathcal{H}$, it follows that $\mathcal{H}$ is the orthogonal direct sum of the $\mathcal{J}$-eigenspaces

$$\mathcal{H}_n := \{v \in \mathcal{H}\colon (\forall t \in \mathcal{J})\ \pi(t)v = t^n v\}.$$

Let π_n denote the corresponding subrepresentation of S on $\mathcal{H}_n$. Accordingly, we write $\iota(v) = \sum_n \iota_n(v)$ with continuous linear maps $\iota_n\colon V \to \mathcal{H}_n$. This leads to the weakly convergent series

$$\varphi(s) = \iota^*\pi(s)\iota = \sum_n \iota_n^*\pi_n(s)\iota_n.$$

We further have $\|\iota_n\| \leq \|\iota\|$ and, for $a \in \mathcal{A}$ with $p(a) < r < 1$,

$$\|\pi_n(a)\| = r^n\|\pi_n(r^{-1}a)\| \leq r^n\|\pi(r^{-1}a)\| \leq r^n.$$

Therefore the above series converges uniformly on every ball B_r, $r < 1$. Further, the homogeneity of the functions $\iota_n^*\pi_n(s)\iota_n$ implies that $\varphi_n(s) = \iota_n^*\pi_n(s)\iota_n$ for every $s \in S$. This in turn implies that $\mathcal{H}_n = \mathcal{H}_{\varphi_n}$. □

5.3 Homogeneous positive definite functions

We now take a closer look at the representations corresponding to homogeneous positive definite functions φ of degree n on $\mathtt{ball}(\mathcal{A}, p)$, resp., $\mathcal{A}$. In particular, we shall see that these functions are completely positive, and this will complete the proof of the result that bounded analytic positive definite functions are completely positive.

Lemma 5.6. *If $(\mathcal{A}_j, p_j)$, $j = 1, 2$, are seminormed involutive algebras, then their projective tensor product $(\mathcal{A}_1 \otimes \mathcal{A}_2, p_1 \otimes p_2)$ with*

$$(p_1 \otimes p_2)(z) := \inf\Big\{ \sum_j p_1(x_j)p_2(y_j) \colon z = \sum_j x_j \otimes y_j \Big\}$$

is also a seminormed involutive algebra.

Proof. It is well-known that $p_1 \otimes p_2$ defines a seminorm on the real associative $*$-algebra $\mathcal{B} := \mathcal{A}_1 \otimes \mathcal{A}_2$ ([Tr67]). For $z = \sum_j x_j \otimes y_j$ and $z' = \sum_j x'_j \otimes y'_j$, we have $zz' = \sum_{j,k} x_j x'_k \otimes y_j y'_k$. From

$$\sum_{j,k} p_1(x_j x'_k)p_2(y_j y'_k) \le \sum_{j,k} p_1(x_j)p_2(y_j)p_1(x'_k)p_2(y'_k) = \sum_j p_1(x_j)p_2(y_j) \sum_k p_1(x'_k)p_2(y'_k)$$

we then derive that

$$(p_1 \otimes p_2)(zz') \le (p_1 \otimes p_2)(z)(p_1 \otimes p_2)(z'),$$

i.e., that $p_1 \otimes q_2$ is submultiplicative. It is also easy to see that $p_1 \otimes p_2$ is involutive. □

Example 5.7. The preceding lemma implies in particular, that we obtain for any seminormed involutive algebra $(\mathcal{A}, p)$ a natural seminorm on the matrix algebra $M_n(\mathcal{A}) \cong M_n(\mathbb{R}) \otimes_{\mathbb{R}} \mathcal{A}$, where the norm on $M_n(\mathbb{R})$ is the operator norm with respect to the euclidean norm on $\mathbb{R}^n$.

We consider the involutive algebra

$$S^n(\mathcal{A}) := (\mathcal{A}^{\otimes n})^{S_n}$$

and endow it with the seminorm p_n obtained by restricting the (projective) seminorm $p^{\otimes n}$ (Lemma 5.6). Then the natural homomorphism $\eta_{\mathcal{A}}^{\otimes n} \colon \mathcal{A}^{\otimes n} \to C^*(\mathcal{A}, p)^{\otimes n}$ intertwines the S_n-actions, so that it induces a morphism

$$S^n(\eta_{\mathcal{A}}) \colon S^n(\mathcal{A}) \to S^n(C^*(\mathcal{A}, p)).$$

If $\mathcal{B}$ and $\mathcal{C}$ are C^*-algebras, we write $\mathcal{B} \otimes \mathcal{C}$ for the C^*-algebra obtained from the algebraic tensor product by completion with respect to the maximal C^*-norm, which leads to the *projective C^*-tensor product* (cf. [Tak02, Def. IV.4.5]).

The following proposition generalizes [Ok66, Thm.3] that deals with the special case of Banach $*$-algebras with approximate identities.

Proposition 5.8. *Let $(\mathcal{A}_j, p_j)$, $j = 1, 2$, be two real seminormed involutive algebras. Then the map $\eta_{\mathcal{A}_1} \otimes \eta_{\mathcal{A}_2} \colon \mathcal{A}_1 \otimes \mathcal{A}_2 \to C^*(\mathcal{A}_1, p_1) \otimes C^*(\mathcal{A}_2, p_2)$ has the universal property of the enveloping C^*-algebra of the seminormed involutive algebra $(\mathcal{A}_1 \otimes \mathcal{A}_2, p_1 \otimes p_2)$.*

Proof. Let $\pi\colon \mathcal{A}_1 \otimes \mathcal{A}_2 \to B(\mathcal{H})$ be a non-degenerate $p_1 \otimes p_2$-bounded representation. If the algebras $\mathcal{A}_j$ are both unital, then $\pi_1(a_1) := \pi(a_1 \otimes \mathbf{1})$ and $\pi_2(a_2) := \pi(\mathbf{1} \otimes a_2)$ are pairwise commuting representations of $\mathcal{A}_1$, resp., $\mathcal{A}_2$ with

$$\pi(a_1 \otimes a_2) = \pi_1(a_1)\pi_2(a_2).$$

If the algebras $\mathcal{A}_j$ are not unital, we have to invoke multiplier techniques to obtain the representations π_j. Let $(\mathcal{A}_j^1, p_j^1)$ denote the unital seminormed involutive algebra associated to $(\mathcal{A}_j, p_j)$ (Definition 2.15). Then $\mathcal{A}_1 \otimes \mathcal{A}_2$ is an ideal of $\mathcal{A}_1^1 \otimes \mathcal{A}_2^1$, so that the latter algebra acts on $\mathcal{A}_1 \otimes \mathcal{A}_2$ by multipliers. For $m_j \in \mathcal{A}_j^1$ and $z \in \mathcal{A}_1 \otimes \mathcal{A}_2$, we have

$$\begin{aligned}
(p_1 \otimes p_2)((m_1 \otimes m_2)z) &= \inf\Big\{ \sum_k p_1(x_k)p_2(y_k)\colon \sum_k x_k \otimes y_k = (m_1 \otimes m_2)z \Big\} \\
&\le \inf\Big\{ \sum_k p_1(m_1x_k)p_2(m_2y_k)\colon \sum_k x_k \otimes y_k = z \Big\} \\
&\le p_1^1(m_1)p_2^1(m_2) \inf\Big\{ \sum_k p_1(x_k)p_2(y_k)\colon \sum_k x_k \otimes y_k = z \Big\} \\
&= p_1^1(m_1)p_2^1(m_2)(p_1 \otimes p_2)(z).
\end{aligned}$$

We conclude that, for $S_j := \overline{\mathtt{ball}}(\mathcal{A}_1^1, p_1^1)$ and $S := \mathtt{ball}(\mathcal{A}_1 \otimes \mathcal{A}_2, p_1 \otimes p_2)$, we have

$$(S_1 \times S_2)S \subseteq S.$$

Applying Lemma 3.22 to the ideal

$$\mathcal{A} := \mathcal{A}_1 \otimes \mathcal{A}_2 \trianglelefteq \mathcal{B} := (\mathcal{A}_1 \otimes \mathcal{A}_2) + \mathcal{A}_1^1 \otimes \mathbf{1},$$

we obtain a unique linear representation $\pi_1^1\colon \mathcal{A}_1^1 \to B(\mathcal{H})$, bounded on S_1 and satisfying

$$\pi_1^1(a)\pi(a_1 \otimes a_2) = \pi(aa_1 \otimes a_2) \quad \text{for} \quad a, a_1 \in \mathcal{A}_1, a_2 \in \mathcal{A}_2.$$

We likewise obtain a unique linear representation $\pi_2^1\colon \mathcal{A}_2^1 \to B(\mathcal{H})$, bounded on S_2 and satisfying

$$\pi_2^1(a)\pi(a_1 \otimes a_2) = \pi(a_1 \otimes aa_2) \quad \text{for} \quad a_1 \in \mathcal{A}_1, a, a_2 \in \mathcal{A}_2.$$

We put $\pi_j := \pi_j^1|_{\mathcal{A}_j}$. For $a_1, a_1' \in \mathcal{A}$ and $a_2, a_2' \in \mathcal{A}_2$, the relation

$$\pi_1(a_1)\pi_2(a_2)\pi(a_1' \otimes a_2') = \pi(a_1a_1' \otimes a_2a_2') = \pi(a_1 \otimes a_2)\pi(a_1' \otimes a_2')$$

now implies that

$$\pi(a_1 \otimes a_2) = \pi_1(a_1)\pi_2(a_2).$$

Since π_j^1 is bounded on S_j, it follows that the representation π_j of $\mathcal{A}_j$ is bounded on $\mathtt{ball}(\mathcal{A}_j, p_j)$, i.e., $\|\pi_j(a_j)\| \le p_j(a_j)$ for $a_j \in \mathcal{A}_j$, $j = 1, 2$. Hence there exist pairwise commuting representations $\widehat{\pi}_j\colon C^*(\mathcal{A}_j, p_j) \to B(\mathcal{H})$ with $\widehat{\pi}_j \circ \eta_{\mathcal{A}_j} = \pi_j$. Now

$$\widehat{\pi}(a_1 \otimes a_2) := \widehat{\pi}_1(a_1)\widehat{\pi}_2(a_2)$$

defines a representation $\widehat{\pi}\colon C^*(\mathcal{A}_1, p_1) \otimes C^*(\mathcal{A}_2, p_2) \to B(\mathcal{H})$ with $\widehat{\pi} \circ (\eta_{\mathcal{A}} \otimes \eta_{\mathcal{A}_2}) = \pi$ (see [Tak02, Prop. IV.4.7]). ∎

Iterating the preceding lemma, we obtain in particular:

Corollary 5.9. *The map $\eta_{\mathcal{A}}^{\otimes n}\colon \mathcal{A}^{\otimes n} \to C^*(\mathcal{A},p)^{\otimes n}$ is universal for the seminormed involutive algebra $(\mathcal{A}^{\otimes n}, p^{\otimes n})$.*

Lemma 5.10. *Let Γ be a finite group acting on the seminormed involutive algebra $(\mathcal{A},p)$. Then the fixed point projection $p\colon \mathcal{A} \to \mathcal{A}^\Gamma$ is completely positive.*

Proof. Since $M_n(p)\colon M_n(\mathcal{A}) \to M_n(\mathcal{A}^\Gamma)$ is the fixed point projection for the canonical action of Γ on $M_n(\mathcal{A})$, it suffices to show that p is positive. This follows from the positivity of

$$p(a^*a) = \frac{1}{|\Gamma|}\sum_{\gamma\in\Gamma}\gamma(a^*a) = \frac{1}{|\Gamma|}\sum_{\gamma\in\Gamma}\gamma(a)^*\gamma(a). \qquad \square$$

Proposition 5.11. *The homomorphism $S^n(\eta_{\mathcal{A}})\colon S^n(\mathcal{A}) \to S^n(C^*(\mathcal{A},p))$ has the universal property of $C^*(S^n(\mathcal{A}),p_n)$.*

Proof. **Case 1:** First we assume that $\mathcal{A}$ is unital. Let $\pi\colon S^n(\mathcal{A}) \to B(\mathcal{H})$ be a p_n-bounded (unital) representation. The S_n-equivariant projection $q\colon \mathcal{A}^{\otimes n} \to S^n(\mathcal{A})$ is completely positive (Lemma 5.10) and since the seminorm $p^{\otimes n}$ is S_n-invariant, q is a $p^{\otimes n}$-contraction. Therefore $\pi\circ q\colon \mathcal{A}^{\otimes n} \to B(\mathcal{H})$ is $p^{\otimes n}$-bounded and completely positive, hence of the form $\iota^*\rho(\cdot)\iota$ for a $p^{\otimes n}$-bounded representation ρ of $\mathcal{A}^{\otimes n}$ (cf. Theorem 3.18). In view of Proposition 3.9 and Corollary 5.9, this implies the existence of a completely positive map

$$\Phi\colon C^*(\mathcal{A},p)^{\otimes n} \to B(\mathcal{H}) \quad \text{with} \quad \Phi(\eta_{\mathcal{A}}(a_1)\otimes\cdots\otimes\eta_{\mathcal{A}}(a_n)) = \pi(q(a_1\otimes\cdots\otimes a_n)).$$

Then $\Phi(\eta_{\mathcal{A}}(a)\otimes\cdots\otimes\eta_{\mathcal{A}}(a)) = \pi(a\otimes\cdots\otimes a))$ implies that

$$\Phi\circ S^n(\eta_{\mathcal{A}}) = \pi.$$

As π is a representation and the image of $S^n(\eta_{\mathcal{A}})$ spans a dense subspace in $S^n(C^*(\mathcal{A},p))$, Φ is multiplicative on $S^n(C^*(\mathcal{A},p))$, so that $\Phi|_{S^n(\mathcal{A})}$ is a representation.

Case 2: If $\mathcal{A}$ is not unital, we assume that π is non-degenerate. We first observe that $S^n(\mathcal{A}) \trianglelefteq S^n(\mathcal{A}^1)$ is an ideal. Let

$$S := \mathtt{ball}(S^n(\mathcal{A}),p_n) \quad \text{and} \quad S^1 := \{b\in S^n(\mathcal{A}^1)\colon bS\cup Sb\subseteq S\}.$$

Then S^1 contains all elements of the form $a^{\otimes n}$, where $a\in\mathtt{ball}(\mathcal{A}^1,p^1)$, and these elements span $S^n(\mathcal{A}^1)$. Therefore Lemma 3.22 implies that π extends to a linear unital representation $\widehat{\pi}\colon S^n(\mathcal{A}^1) \to B(\mathcal{H})$ which is bounded on S^1.

We claim that $\widehat{\pi}$ is bounded on $\mathtt{ball}(S^n(\mathcal{A}^1),p_n^1)$. To this end, we first observe that, for $a\in\mathcal{A}$, $m\in\mathcal{A}^1$, we have $p(ma)\le p^1(m)p(a)$ because p^1 is submultiplicative and extends p. This leads to

$$p^{\otimes n}((m_1\otimes\cdots\otimes m_n)z) \le p^1(m_1)\cdots p^1(m_n)p^{\otimes n}(z) \quad \text{for} \quad m_j\in\mathcal{A}^1, z\in\mathcal{A}^{\otimes n},$$

and this in turn to

$$p^{\otimes n}(wz) \le (p^1)^{\otimes n}(w)p^{\otimes n}(z) \quad \text{for} \quad w\in(\mathcal{A}^1)^{\otimes n}, z\in\mathcal{A}^{\otimes n}.$$

This specializes to

$$p_n(wz) \leq p_n^1(w)p_n(z) \quad \text{for} \quad w \in S^n(\mathcal{A}^1), z \in S^n(\mathcal{A}).$$

From this estimate we derive that $\mathtt{ball}(S^n(\mathcal{A}^1), p_n^1) \subseteq S^1$, so that $\widehat{\pi}$ is bounded on the semigroup $\mathtt{ball}(S^n(\mathcal{A}^1), p_n^1)$, which leads to $\|\widehat{\pi}(w)\| \leq p_n^1(w)$ for $w \in S^n(\mathcal{A}^1)$. Hence Case 1 shows that there exists a unital representation $\beta\colon S^n(C^*(\mathcal{A}^1, p)) \to B(\mathcal{H})$ with

$$\beta \circ \eta_{S^n(\mathcal{A}^1)} = \widehat{\pi}.$$

Composing with the natural map $S^n(C^*(\mathcal{A}, p)) \to S^n(C^*(\mathcal{A}, p)^1) \cong S^n(C^*(\mathcal{A}^1, p^1))$ (cf. Remark 2.16), we obtain a representation $\gamma\colon S^n(C^*(\mathcal{A}, p)) \to B(\mathcal{H})$ with

$$\gamma(S^n(\eta_{\mathcal{A}})(a^{\otimes n})) = \gamma(\eta_{\mathcal{A}}(a)^{\otimes n}) = \beta(\eta_{\mathcal{A}^1}(a)^{\otimes n}) = \beta(\eta_{S^n(\mathcal{A}^1)}(a^{\otimes n})) = \widehat{\pi}(a^{\otimes n}) = \pi(a^{\otimes n}).$$

This implies that the two representations π and $\gamma \circ S^n(\eta_{\mathcal{A}})$ of $S^n(\mathcal{A})$ coincide, and thus $S^n(\eta_{\mathcal{A}})\colon S^n(\mathcal{A}) \to S^n(C^*(\mathcal{A}, p))$ has the universal property of the enveloping C^*-algebra of the seminormed involutive algebra $(S^n(\mathcal{A}), p_n)$. □

The following automatic continuity lemma is a version of Corollary 3.9 for polynomials of higher degree. We recall that the dilatability assumption is always satisfied if $\mathcal{A}$ is unital (Theorem 3.18).

Lemma 5.12. *Let $(\mathcal{A}, p)$ be a real seminormed involutive algebra, $n \in \mathbb{N}_0$, and V be a complex Hilbert space. For every p-bounded dilatable positive definite homogeneous polynomial $\varphi\colon \mathcal{A} \to B(V)$ of degree n, there exists a unique complex linear completely positive map*

$$\Phi_n\colon S^n(C^*(\mathcal{A}, p)) \to B(V) \quad \textit{with} \quad \varphi(a) = \Phi_n(\eta_{\mathcal{A}}(a)^{\otimes n}) \quad \textit{for} \quad a \in \mathcal{A}.$$

In particular, φ is completely positive.

Proof. Let $(\pi, \mathcal{H}, \iota)$ be a dilation of $\varphi|_{\mathtt{ball}(\mathcal{A},p)}$. The assumption that φ is a homogeneous polynomial of degree n implies that all elements of $\mathcal{H} \cong \mathcal{H}_\varphi$ have this property, so that $\pi(ra) := r^n\pi(a)$ yields an extension to a representation of the involutive semigroup $(\mathcal{A}, \cdot)$. The map $\pi\colon \mathcal{A} \to B(\mathcal{H})$ is a polynomial of degree n bounded on $\mathtt{ball}(\mathcal{A}, p)$. Hence π defines a unique linear representation $\widetilde{\pi}\colon S^n(\mathcal{A}) \to B(\mathcal{H})$ bounded on the subsemigroup

$$S := \{a^{\otimes n}\colon a \in \mathtt{ball}(\mathcal{A}, p)\} \subseteq \mathtt{ball}(S^n(\mathcal{A}), p_n).$$

This implies that the corresponding symmetric n-linear map $\beta\colon \mathcal{A}^n \to B(\mathcal{H})$ is continuous on the diagonal with respect to the p-topology, and polarization shows that β is continuous with respect to the p-topology on $\mathcal{A}^n$. This in turn shows that the corresponding linear map $\widetilde{\beta}\colon \mathcal{A}^{\otimes n} \to B(\mathcal{H})$ is $p^{\otimes n}$-continuous, and therefore $\widetilde{\beta}|_{S^n(\mathcal{A})} = \widetilde{\pi}$ is p_n-continuous, i.e., bounded on $\mathtt{ball}(S^n(\mathcal{A}), p_n)$, and therefore $\|\widetilde{\pi}(z)\| \leq 1$ for $z \in \mathtt{ball}(S^n(\mathcal{A}), p_n)$ (cf. Remark 2.4).

Now Proposition 5.11 provides a representation $\rho\colon S^n(C^*(\mathcal{A}, p)) \to B(\mathcal{H})$ with $\rho \circ S^n(\eta_{\mathcal{A}}) = \widetilde{\pi}$. We thus obtain for $a \in \mathcal{A}$:

$$\varphi(a) = \iota^*\pi(a)\iota = \iota^*\widetilde{\pi}(a^{\otimes n})\iota = \iota^*\rho(\eta_{\mathcal{A}}(a)^{\otimes n})\iota.$$

Hence $\varphi = \Phi_n(\eta_{\mathcal{A}}(a)^{\otimes n})$ holds for $\Phi_n(b) := \iota^*\rho(b)\iota$, which is obviously completely positive, hence also positive definite.

For $\Phi(b) := \Phi_n(b^{\otimes n})$ we now have $\Phi \circ \eta_{\mathcal{A}} = \varphi$. Therefore the complete positivity of φ follows from the complete positivity of $\eta_{\mathcal{A}}$, the complete positivity of the map $C^*(\mathcal{A}, p) \to S^n(C^*(\mathcal{A}, p)), b \mapsto b^{\otimes n}$ ([Arv87, Lemma 3.5]) and the complete positivity of Φ_n. □

Theorem 5.13. *If $(\mathcal{A}, p)$ is a seminormed algebra, V a complex Hilbert space and $\varphi\colon \mathtt{ball}(\mathcal{A}, p) \to B(V)$ a dilatable bounded positive definite function which is analytic with respect to some locally convex topology for which p is continuous, then φ is completely positive.*

Proof. First, Lemma 5.5 asserts that the Taylor series $\varphi = \sum_n \varphi_n$ converges uniformly on all balls B_r, $r < 1$. Next, Lemma 5.12 implies that the functions φ_n are completely positive, so that φ is completely positive as well. □

6 Proof of the main theorem

In this section we finally prove Theorem 1.1.

Definition 6.1. For any C^*-algebra $\mathcal{B}$ and $k \in \mathbb{N}_0$, we consider the C^*-algebra

$$S^n(\mathcal{B}) := (\mathcal{B}^{\otimes n})^{S_n}$$

generated by the elements of the form $b^{\otimes n}$, $b \in \mathcal{B}$, in the C^*-algebra $\mathcal{B}^{\otimes n}$. Here the tensor product means the *projective C^*-tensor product*, resp., the completion of the tensor product with respect to the maximal C^*-norm (cf. [Tak02, Def. IV.4.5]). The above $S^n(\mathcal{B})$ is the subalgebra of fixed points for the natural action of the symmetric group S_n on $\mathcal{B}^{\otimes n}$. For $n = 0$, we put $S^0(\mathcal{B}) := \mathbb{C}$. The *exponential C^*-algebra* of $\mathcal{B}$ is

$$e^{\mathcal{B}} := \Big\{(b_k)_{k\geq 0} \in \prod_{k\geq 0} S^k(\mathcal{B}) \mid \lim_{k\to\infty} \|b_k\| = 0\Big\}. \tag{10}$$

We then obtain a natural map

$$\Gamma_{\mathcal{B}}\colon \mathtt{ball}(\mathcal{B}) \to \overline{\mathtt{ball}}(e^{\mathcal{B}}), \quad \Gamma_{\mathcal{B}}(b) = \sum_{k\geq 0} b^{\otimes k}.$$

We will write simply $\Gamma_{\mathcal{B}} = \Gamma$ whenever the C^*-algebra $\mathcal{B}$ is understood from the context.

Lemma 6.2. *For every C^*-algebra $\mathcal{B}$, the map $\Gamma_{\mathcal{B}}\colon \mathtt{ball}(\mathcal{B}) \to C(e^{\mathcal{B}})$ is a completely positive holomorphic homomorphism of $*$-semigroups.*

Proof. For every $k \in \mathbb{N}_0$, the map $\Gamma_k|_{\mathtt{ball}(\mathcal{B})}\colon \mathtt{ball}(\mathcal{B}) \to S^k(\mathcal{B})$ is completely positive and homogeneous of degree k by [Arv87, Lemma 3.5]. Since their sum converges uniformly on every ball of radius $r < 1$, the map $\Gamma_{\mathcal{B}}$ is also holomorphic and completely positive. □

We now consider the natural map

$$\Gamma = \Gamma_{\mathcal{A}}\colon \mathtt{ball}(\mathcal{A}, p) \to \mathtt{ball}(e^{C^*(\mathcal{A},p)}), \quad \Gamma(a) = \sum_{k\geq 0} \eta_{\mathcal{A}}(a)^{\otimes k}.$$

Lemma 6.3. *The map* $\Gamma\colon \mathtt{ball}(\mathcal{A},p) \to e^{C^*(\mathcal{A},p)}$ *is completely positive and analytic in the* p*-topology on* $\mathcal{A}$*.*

Proof. Let $\mathcal{B} := C^*(\mathcal{A},p)$. Since the map $\eta_{\mathcal{A}}\colon \mathtt{ball}(\mathcal{A},p) \to \mathtt{ball}(\mathcal{B})$ is continuous linear, the analyticity of Γ follows from the holomorphy of $\Gamma_{\mathcal{B}}$ (Lemma 6.2).

To see that Γ is completely positive, we have to show that, for $A = (a_{ij}) \geq 0$ in $M_n(\mathcal{A})$ and $p(a_{ij}) < 1$, the matrix $(\Gamma(a_{ij})) \in M_n(e^{\mathcal{B}})$ is positive. This follows from the complete positivity of the linear $*$-homomorphism $\eta_{\mathcal{A}}\colon \mathcal{A} \hookrightarrow \mathcal{B}$ (Remark 2.10(b)) and Lemma 6.2. □

The following theorem is our main theorem stated as Theorem 1.1 in the introduction.

Theorem 6.4. *Let* $(\mathcal{A},p)$ *be a real seminormed involutive algebra,* V *a complex Hilbert space and* $\varphi\colon \mathtt{ball}(\mathcal{A},p) \to B(V)$ *be a bounded function. Then the following are equivalent:*

(i) φ *is completely positive and dilatable.*

(ii) φ *is dilatable, positive definite and analytic with respect to some locally convex topology for which* p *is continuous.*

(iii) *There exists a linear completely positive map* $\Phi\colon e^{C^*(\mathcal{A},p)} \to B(V)$ *with* $\Phi \circ \Gamma = \varphi$*.*

Proof. (i) ⇒ (ii) follows from Theorem 4.12.

(ii) ⇒ (i) follows from Theorem 5.13.

(iii) ⇒ (i): Let $\mathcal{B} := C^*(\mathcal{A},p)$ and $\Phi\colon e^{\mathcal{B}} \to B(V)$ be any completely positive linear map. Since a composition of completely positive maps is completely positive (Remark 2.10), it follows from Lemma 6.3 that $\Phi \circ \Gamma$ is an analytic completely positive map. Further, the dilatability of φ follows from the dilatability of Φ (cf. Proposition 3.9).

(i) ⇒ (iii): First, Theorem 4.12 implies the existence of an expansion $\varphi = \sum_{n=0}^{\infty} \varphi_n$, where each φ_n is a homogeneous completely positive polynomial with $\|\varphi_n\|_\infty \leq \|\varphi\|_\infty$. Lemma 5.12 now provides linear positive maps $\Phi_n\colon S^n(C^*(\mathcal{A},p)) \to B(V)$ with $\varphi_n(a) = \Phi_n(\eta_{\mathcal{A}}(a)^{\otimes n})$ for $a \in \mathtt{ball}(\mathcal{A},p)$. For the canonical extensions to $\mathcal{A}^1$, we further have $\Phi_n^1(\mathbf{1}^{\otimes n}) = \varphi_n^1(\mathbf{1}) \leq \varphi^1(\mathbf{1})$, so that the sequence $(\|\Phi_n^1\|)_{n\in\mathbb{N}_0}$ is bounded. This implies that $\Phi(x) := \sum_n \Phi_n(x_n)$ defines a continuous linear function $e^{C^*(\mathcal{A},p)} \to B(V)$ with the required properties. □

For unital algebras, we obtain the following simplification:

Corollary 6.5. *Let* $(\mathcal{A},p)$ *be a real unital seminormed involutive algebra,* V *a complex Hilbert space and* $\varphi\colon \mathtt{ball}(\mathcal{A},p) \to B(V)$ *be a bounded function. Then the following are equivalent:*

(i) φ *is completely positive.*

(ii) φ *is positive definite and analytic with respect to some locally convex topology for which* p *is continuous.*

(iii) *There exists a linear completely positive map* $\Phi\colon e^{C^*(\mathcal{A},p)} \to B(V)$ *with* $\Phi \circ \Gamma = \varphi$*.*

Proof. If $\mathcal{A}$ is unital, then every bounded positive definite function on $\mathtt{ball}(\mathcal{A},p)$ is dilatable by Theorem 3.18, so that the assertion follows from Theorem 6.4. □

Recall from Example 3.10 that the dilatability assumptions cannot be removed from Theorem 6.4, which is the same as Theorem 1.1.

Corollary 6.6. *The assignment $\Phi \mapsto \Phi \circ \Gamma$ is a bijection between the completely positive linear $B(V)$-valued maps on the C^*-algebra $e^{C^*(\mathcal{A},p)}$ and the bounded dilatable completely positive $B(V)$-valued functions on the $*$-semigroup $\mathtt{ball}(\mathcal{A},p)$.*

Proof. That $\Phi \circ \Gamma$ determines Φ uniquely follows from the fact that the complex linear span of $\Gamma(\mathtt{ball}(\mathcal{A},p))$ is dense in $e^{C^*(\mathcal{A},p)}$ ([Arv87, Lemma 3.23]). □

Remark 6.7. If the real seminormed involutive algebra $(\mathcal{A},p) = (\mathcal{A}, \|\cdot\|)$ is a unital C^*-algebra, then $C^*(\mathcal{A},p) \cong \mathcal{A}_{\mathbb{C}} \cong \mathcal{A} \oplus \overline{\mathcal{A}}$ and the completely positive linear maps on $\Phi\colon e^{\mathcal{A}\oplus\overline{\mathcal{A}}} \cong e^{\mathcal{A}} \otimes e^{\overline{\mathcal{A}}} \to \mathbb{C}$ normalized by $\|\Phi\| = 1$ are precisely the states of $e^{\mathcal{A}} \otimes e^{\overline{\mathcal{A}}}$. This special case of the above Theorem 6.4 is [Arv87, Th. 3.1].

The following theorem is an important consequence of Theorem 6.4 for the representation theory of $\mathtt{ball}(\mathcal{A},p)$.

Theorem 6.8. *Let $(\mathcal{A},p)$ be a real seminormed involutive algebra and $\mathcal{H}$ a complex Hilbert space. Then the correspondence $\widehat{\pi} \mapsto \widehat{\pi} \circ \Gamma$ defines a commutant-preserving bijection between the $*$-representations of the C^*-algebra $e^{C^*(\mathcal{A},p)}$ on $\mathcal{H}$ and the bounded real-analytic $*$-representations of the $*$-semigroup $\mathtt{ball}(\mathcal{A},p)$ on $\mathcal{H}$.*

Proof. Every $*$-representation of a $*$-semigroup is a dilatable positive definite function, hence we can use Theorem 6.4 to see that a bounded real-analytic unital $*$-representation $\pi\colon \mathtt{ball}(\mathcal{A}) \to B(\mathcal{H})$ is completely positive and that there exists a unique completely positive linear map $\Phi\colon e^{C^*(\mathcal{A},p)} \to B(\mathcal{H})$ with $\pi = \Phi \circ \Gamma$. It remains to prove that, for all $x, y \in e^{C^*(\mathcal{A},p)}$, we have $\Phi(xy) = \Phi(x)\Phi(y)$. Since Φ is positive, it is also continuous even though it is defined on the non-unital C^*-algebra $e^{C^*(\mathcal{A},p)}$ (see [Pau02, Ch. 2]). As $\pi = \Phi \circ \Gamma$ and both π and Γ are semigroup homomorphisms, $\Phi(xy) = \Phi(x)\Phi(y)$ holds for $x, y \in \Gamma(\mathtt{ball}(\mathcal{A},p))$, and since $\Gamma(\mathtt{ball}(\mathcal{A},p))$ spans a dense linear subspace of $e^{C^*(\mathcal{A},p)}$ and Φ is continuous, it is also multiplicative, hence a $*$-representation of the C^*-algebra $e^{C^*(\mathcal{A},p)}$.

Conversely, if Φ is a $*$-homomorphism of C^*-algebras, then $\Phi\circ\Gamma$ is a bounded real-analytic $*$-representation of the $*$-semigroup $\mathtt{ball}(\mathcal{A},p)$ on $\mathcal{H}$. □

Remark 6.9. (a) Theorem 6.8 asserts that the bounded real-analytic representation theory of the involutive semigroup $\mathtt{ball}(\mathcal{A},p)$ is faithfully represented by the C^*-algebra $e^{C^*(\mathcal{A},p)}$, so this is a host algebra in the sense of [Gr05]. Since the real linear map $\eta_{\mathcal{A}}\colon \mathcal{A} \to C^*(\mathcal{A},p)$ extends to a complex linear map $\mathcal{A}_{\mathbb{C}} \to C^*(\mathcal{A},p)$, it follows in particular that every bounded analytic completely positive function $\varphi\colon \mathtt{ball}(\mathcal{A},p) \to B(V)$ extends to a holomorphic function on the open semigroup $\mathtt{ball}(\mathcal{A}_{\mathbb{C}}, p_{\mathbb{C}})$ for $p_{\mathbb{C}}(a+ib) := p(a) + p(b)$ (cf. Definition 2.12). Therefore $e^{C^*(\mathcal{A},p)}$ is also a host algebra for the bounded holomorphic representations theory of the complex semigroup $\mathtt{ball}(\mathcal{A}_{\mathbb{C}}, q)$ in the sense of [Ne08].

(b) Since $e^{C^*(\mathcal{A},p)}$ is a c_0-direct sum of the ideals $S^N(C^*(\mathcal{A},p))$, all its non-degenerate representations decompose as direct sums of representations on which exactly one of these ideals acts non-trivially. Therefore the main point in applying Theorem 6.8 is to identify the representations of $S^N(C^*(\mathcal{A},p)) \cong C^*(S^N(\mathcal{A}), p_N)$ in terms of the (linear) representations of the C^*-algebra $C^*(\mathcal{A},p)$, resp., representations of the algebra $\mathcal{A}$ which are bounded on $\mathtt{ball}(\mathcal{A},p)$.

(c) For every $N \in \mathbb{N}$, we see that C^*-algebra $S^N(C^*(\mathcal{A},p))$ is a host algebra for the class of the bounded N-homogeneous polynomial representations of $\mathtt{ball}(\mathcal{A},p)$, resp., the N-homogeneous polynomial representations of $\mathcal{A}$ bounded on $\mathtt{ball}(\mathcal{A},p)$.

Corollary 6.10. *Let $\mathcal{A}$ be a C^*-algebra, considered as a real seminormed algebra, and $\mathcal{H}$ be a complex Hilbert space. Every bounded real-analytic $*$-representation $\pi\colon \mathtt{ball}(\mathcal{A}) \to B(\mathcal{H})$ is completely positive and the correspondence $\Phi \mapsto \Phi \circ (\Gamma_{\mathcal{A}} \otimes \Gamma_{\overline{\mathcal{A}}})$ is a bijection between the $*$-representations of the C^*-algebra $e^{\mathcal{A}} \otimes e^{\overline{\mathcal{A}}}$ on $\mathcal{H}$ and the bounded real-analytic $*$-representations of the $*$-semigroup $\mathtt{ball}(\mathcal{A})$ on $\mathcal{H}$.*

Proof. This is the special case of Theorem 6.8, where $\mathcal{A}$ is a unital C^*-algebra. □

Example 6.11. (cf. [HN87, Ex. 2.1]) We consider the non-unital C^*-algebra $\mathcal{A} = c_0(\mathbb{N}_0, \mathbb{C})$. Then

$$\varphi(z) := \sum_{n=0}^{\infty} z_n^n$$

is a completely positive $\mathbb{C}$-valued function on $\mathcal{A}$ which is bounded on the ball $B_s(0)$ of radius s if and only if $s < 1$. In particular, it is unbounded on $\mathtt{ball}(\mathcal{A})$. For $r > 0$ we put $\varphi_r(z) := \varphi(rz)$. Then φ_r is bounded on $\mathtt{ball}(\mathcal{A})$ if $r < 1$, hence dilatable because $\mathcal{A}$ is a C^*-algebra (Proposition 3.9).

In view of

$$\varphi(zw^*) := \sum_{n=0}^{\infty} z_n^n \overline{w_n}^n,$$

the functions $\varphi_n(z) = z_n^n$ form an orthonormal basis of the Hilbert space $\mathcal{H}_\varphi$, which implies in particlar that $\varphi \notin \mathcal{H}_\varphi$. For $r < 1$, we have $\|\varphi_n\| = r^{-n/2}$, so that $\sum_n r^{2n}\|\chi_n\|^2 = \sum_n r^n < \infty$ implies that $\varphi_r \in \mathcal{H}_{\varphi_r}$ holds for $r < 1$ (cf. Proposition 3.7).

7 Relations to Arveson's c.p. concept

Let $\mathcal{A}$ and $\mathcal{B}$ be C^*-algebras. We say that a map $\varphi\colon \mathtt{ball}(\mathcal{A}) \to \mathcal{B}$ is *completely positive of type (W)* if every positive element $A = (a_{ij}) \in M_n(\mathcal{A})$ with $\|A\| < 1$ is mapped to a positive element of $\mathcal{B}$. This is the concept of complete positivity used in [Arv87] for functions on the open unit ball of unital a C^*-algebra.

Remark 7.1. (a) If $\varphi\colon \mathtt{ball}(\mathcal{A}) \to \mathcal{B}$ is homogeneous of degree $\alpha \in \mathbb{R}$ as in Lemma 2.11, then complete positivity of type (W) implies complete positivity of its extension $\widehat{\varphi}\colon \mathcal{A} \to \mathcal{B}$. In fact, for any $A = (a_{ij}) \in M_n(\mathcal{A})_+$, there exists an $r > 0$ with $r\|a_{ij}\| < 1$ for all i, j and

$$\|(ra_{ij})\| = r\|(a_{ij})\| < 1.$$

Then $(\widehat{\varphi}(a_{ij})) = r^{-\alpha}(\varphi(ra_{ij})) \in M_n(\mathcal{B})_+$ implies that $\widehat{\varphi}\colon \mathcal{A} \to \mathcal{B}$ is completely positive.

(b) We will show in Example 7.4 that complete positivity of type (W) does not imply positive definiteness. This is related to the fact that the matrix A used in Lemma 2.9 need not be a contraction. For example the matrix

$$A := \begin{pmatrix} r^2 & rs \\ rs & s^2 \end{pmatrix}, \quad 0 < r, s < 1$$

satisfies $\|A\| = r^2 + s^2$.

The next proposition shows that, for bounded positive definite functions on $\mathtt{ball}(\mathcal{A}, p)$, complete positivity of type (W) implies complete positivity in the sense of Definition 2.7(b).

Proposition 7.2. *Let $\mathcal{A}$ be a C^*-algebra and V be a complex Hilbert space. If a bounded positive definite function $\varphi\colon \mathtt{ball}(\mathcal{A}) \to B(V)$ is completely positive of type* (W)*, then it is also completely positive.*

Proof. Let $\varphi\colon \mathtt{ball}(\mathcal{A}) \to B(V)$ be completely positive of type (W). It follows by Remark 7.1(c) that the corresponding GNS representation $(\pi_\varphi, \mathcal{H}_\varphi)$ is completely positive of type (W). Since this representation is non-degenerate, it is a direct sum of cyclic representation $(\pi_j, \mathcal{H}_j)_{j\in J}$ generated by cyclic vectors $v_j \in \mathcal{H}_j \subseteq \mathcal{H}_\varphi$ ([Ne00]). Then the functions

$$\varphi_j(s) := \langle \pi_j(s)v_j, v_j\rangle = \langle \pi(s)v_j, v_j\rangle$$

are also completely positive of type (W) since the linear functional $B(\mathcal{H}_\varphi) \to \mathbb{C}, A \mapsto \langle Av_j, v_j\rangle$ is positive definite, hence completely positive (Remark 2.10(c)). Now Remark 7.1(d) implies that the functions φ_j are completely positive, so that the representations π_j are completely positive by Proposition 4.1. This in turn implies that π_φ is completely positive, which, again by Proposition 4.1, implies that φ is completely positive. □

We will show by a simple example that complete positivity of type (W) does not imply positive definiteness, hence also not complete positivity (see Example 7.4). The following simple observation is the key to that example.

Lemma 7.3. *Let V be any Hilbert space and $A \in B(V)$. If $0 \le A \le \mathbf{1}$ and $u, v \in V$ with $\|u\| = \|v\| = 1$ and $\langle u, v\rangle = 0$, then $|\langle Au, v\rangle| \le 1/2$.*

Proof. Since $A \ge 0$, by the Cauchy–Schwartz inequality we obtain

$$|\langle Au, v\rangle|^2 \le \langle Au, u\rangle\langle Av, v\rangle.$$

Similarly, since $\mathbf{1} - A \ge 0$, we have $|\langle(\mathbf{1} - A)u, v\rangle|^2 \le \langle(\mathbf{1} - A)u, u\rangle\langle(\mathbf{1} - A)v, v\rangle$, and then, by using the hypothesis on u and v, we obtain

$$|\langle Au, v\rangle|^2 \le (1 - \langle Au, u\rangle)(1 - \langle Av, v\rangle).$$

By multiplying the above displayed inequalities and using the fact that $t(1 - t) \le 1/4$ if $0 \le t \le 1$, we obtain $|\langle Au, v\rangle|^4 \le 1/16$, which leads to the asserted inequality. □

Example 7.4. Let $\varphi\colon \mathbb{D} \to \mathbb{C}$ satisfying $\varphi(z) = z$ if $|z| \le 1/2$ or $z \in \mathbb{D} \cap [0, \infty)$. Then the following assertions hold:

1. For every integer $n \ge 1$ and $0 \le A = (a_{ij}) \in M_n(\mathbb{C})$ with $\|A\| < 1$ we have $0 \le (\varphi(a_{ij})) \in M_n(\mathbb{C})$.

2. If there exists $z_0 \in \mathbb{D}$ with $\varphi(\overline{z_0}) \ne \overline{\varphi(z_0)}$, then there exist $a_1, a_2 \in \mathbb{D}$ for which the matrix $(\varphi(a_i\overline{a_j}))_{1\le i,j\le 2} \in M_2(\mathbb{C})$ fails to be nonnegative.

For the first assertion, if $0 \leq A = (a_{ij}) \in M_n(\mathbb{C})$ with $\|A\| < 1$, then $0 \leq A \leq \mathbf{1}$, hence by Lemma 7.3 we obtain $|a_{ij}| \leq 1/2$ if $i \neq j$. Moreover, $0 \leq a_{jj} \leq 1$. Consequently $\varphi(a_{ij}) = a_{ij}$ for all $i, j \in \{1, \ldots, n\}$, and then trivially $0 \leq (\varphi(a_{ij})) \in M_n(\mathbb{C})$.

For the second assertion, select any $a_1, a_2 \in \mathbb{D}$ with $a_1\overline{a_2} = z_0$. Since $\varphi(\overline{z_0}) \neq \overline{\varphi(z_0)}$, it follows that the matrix $(\varphi(a_i\overline{a_j})) \in M_2(\mathbb{C})$ fails to be hermitian, hence in particular it cannot be nonnegative.

The above example shows that [Arv87, Lemma 5.3] actually needs a stronger notion of complete positivity, which should ensure that every completely positive function in the corresponding sense is positive definite, so that [Arv87, Th. 2.2] is applicable in that setting. As we have seen in the present paper, the stronger notion of complete positivity from Definition 2.7(b) not only fills that gap from [Arv87] but also is flexible enough to allow one to develop dilation theory for maps defined on the unit ball of any real seminormed involutive algebra. As explained in the introduction, this level of generality is necessary for the applications to mapping algebras $\mathcal{A} = C^\infty(X, \mathcal{B})$ for any C^*-algebra $\mathcal{B}$, where X is a smooth manifold (see [BN15] for details).

A Analytic functions on infinite dimensional spaces

In this appendix, we collect the definitions that are relevant to say what analytic functions between open subsets of locally convex spaces are.

Definition A.1. (a) Let E and F be locally convex spaces, $U \subseteq E$ open and $f\colon U \to F$ a map. Then the *derivative of f at x in the direction h* is defined as

$$\mathtt{d}f(x)(h) := (\partial_h f)(x) := \lim_{t \to 0} \frac{1}{t}(f(x + th) - f(x))$$

whenever it exists. The function f is called *differentiable at x* if $\mathtt{d}f(x)(h)$ exists for all $h \in E$. It is called *continuously differentiable*, if it is differentiable at all points of U and

$$\mathtt{d}f\colon U \times E \to F, \quad (x, h) \mapsto \mathtt{d}f(x)(h)$$

is a continuous map. Note that this implies that the maps $\mathtt{d}f(x)$ are linear (cf. [GN15, Lemma 2.2.14]). The map f is called a *$\mathcal{C}^k$-map*, $k \in \mathbb{N} \cup \{\infty\}$, if it is continuous, the iterated directional derivatives

$$\mathtt{d}^j f(x)(h_1, \ldots, h_j) := (\partial_{h_j} \cdots \partial_{h_1} f)(x)$$

exist for all integers $1 \leq j \leq k$, $x \in U$ and $h_1, \ldots, h_j \in E$, and all maps $\mathtt{d}^j f\colon U \times E^j \to F$ are continuous. As usual, $\mathcal{C}^\infty$-maps are called *smooth*.

(b) If E and F are complex locally convex spaces, then f is called *complex analytic* if it is continuous and, for each $x \in U$, there exists a 0-neighborhood V with $x + V \subseteq U$ and continuous homogeneous polynomials $\beta_k\colon E \to F$ of degree k such that, for each $h \in V$, we have $f(x+h) = \sum_{k=0}^\infty \beta_k(h)$, as a pointwise limit ([BS71]). The map f is called *holomorphic* if it is $\mathcal{C}^1$ and, for each $x \in U$, the map $\mathtt{d}f(x)\colon E \to F$ is complex linear. If F is sequentially complete, then f is holomorphic if and only if it is complex analytic ([BS71, Ths. 3.1, 6.4]).

(c) If E and F are real locally convex spaces, then we call a map $f\colon U \to F$, $U \subseteq E$ open, *real analytic* or a $\mathcal{C}^\omega$-map, if for each point $x \in U$, there exists an open neighborhood

$V \subseteq E_{\mathbb{C}}$ and a holomorphic map $f_{\mathbb{C}}\colon V \to F_{\mathbb{C}}$ with $f_{\mathbb{C}}|_{U\cap V} = f|_{U\cap V}$. The advantage of this definition, which differs from the one in [BS71], is that it also works nicely for non-complete spaces. Any analytic map is smooth, and the corresponding chain rule holds without any condition on the underlying spaces (see [Gl02] for details).

Acknowledgments: D. Beltiţă was supported by the Grant of the Romanian National Authority for Scientific Research, CNCS-UEFISCDI, project number PN-II-ID-PCE-2011-3-0131, and the Emerging Fields Project "Quantum Geometry" of the University of Erlangen, and K.-H. Neeb was supported by DFG-grant NE 413/7-2, Schwerpunktprogramm "Darstellungstheorie".

References

[AC86] Ando, T., and M.D. Choi, *Nonlinear completely positive maps*, in: R. Nagel, U. Schlotterbeck, M.P.H. Wolff (eds.), "Aspects of Positivity in Functional Analysis", North-Holland Math. Stud., 122, North-Holland, Amsterdam, 1986, pp. 3–13.

[Arv69] Arveson, W., *Subalgebras of C^*-algebras*, Acta Math. **123** (1969), 141–224

[Arv87] —, *Nonlinear states on C^*-algebras*, in: P.E.T. Jorgensen and P.S. Muhly (eds.), "Operator algebras and mathematical physics", Contemp. Math., 62, Amer. Math. Soc., Providence, RI, 1987, pp. 283–343.

[Arv10] —, *Dilation theory yesterday and today*, in S. Axler, P. Rosenthal and D. Sarason (eds.), "A glimpse at Hilbert space operators", Oper. Theory Adv. Appl., 207, Birkhäuser Verlag, Basel, 2010, pp. 99–123.

[BN12] Beltiţă, D., and K.-H. Neeb, *Schur-Weyl theory for C^*-algebras*, Math. Nachr. **285** (2012), no. 10, 1170–1198.

[BN15] Beltiţă, D., and K.-H. Neeb, *Non-linear representations of C^*-algebras and their applications*, in preparation.

[Bha98] Bhatt, S. J., *Stinespring representability and Kadison's Schwarz inequality in non-unital Banach star algebras and applications*, Proc. Indian Acad. Sci. Math. Sci. **108** (1998), no. 3, 283–303.

[Bi10] Biller, H., *Continuous inverse algebras with involution*, Forum Math. **22** (2010), no. 6, 1033–1059.

[BS71] Bochnak, J., and J. Siciak, *Polynomials and multilinear mappings in topological vector spaces*, Studia Math. **39** (1971), 59–76.

[CR78] Christensen, J. P. R., and P. Ressel, *Functions operating on positive definite matrices and a theorem of Schoenberg*, Trans. Amer. Math. Soc. **243** (1978), 89–95.

[Dix64] Dixmier, J., "Les C^*-algèbres et leurs représentations," Gauthier-Villars, Paris, 1964.

[EI13] Enomoto, T., and M. Izumi, *Indecomposable characters of infinite dimensional groups associated with operator algebras*, Preprint, arXiv.OA:1308.6329

[FD88] Fell, J. M. G., and R. S. Doran, "Representations of $*$-Algebras, Locally Compact Groups, and Banach-$*$-Algebraic Bundles I,II," Academic Press, 1988

[Gl02] Glöckner, H., *Infinite-dimensional Lie groups without completeness restrictions*, in: A. Strasburger, J. Hilgert, K.-H. Neeb and W. Wojtyński (eds.), "Geometry and Analysis on Finite- and Infinite-dimensional Lie Groups", Banach Center Publ., 55, Polish Acad. Sci., Warsaw, 2002, pp. 43–59.

[GN15] Glöckner, H., and K.-H. Neeb, "Infinite dimensional Lie groups, Vol. I, Basic Theory and Main Examples," book in preparation

[Gr05] Grundling, H., *Generalizing group algebras*, J. London Math. Soc. **72** (2005), 742–762. An erratum is in J. London Math. Soc. **77** (2008), 270–271

[HN87] Hiai, F., and Y. Nakamura, *Extensions of nonlinear completely positive maps*, J. Math. Soc. Japan **39** (1987), no. 3, 367–384.

[Ho69] Horn, R. A., *The theory of infinitely divisible matrices and kernels*, Trans. Amer. Math. Soc. **136** (1969), 269–286

[JN13] Janssens, B., and K. H. Neeb, *Norm continuous unitary representations of Lie algebras of smooth sections*, Internat. Math. Res. Notices, to appear; arXiv:math.RT:1302.2535

[Jo64] Johnson, B. E., *An introduction to the theory of centralizers*, Proc. London Math. Soc. (3) **14** (1964), 299–320

[Ki73] Kirillov, A. A., *Representation of the infinite-dimensional unitary group*, Dokl. Akad. Nauk. SSSR **212**(1973), 288–290.

[Ne00] Neeb, K.-H., "Holomorphy and Convexity in Lie Theory". de Gruyter Expositions in Mathematics **28**, Walter de Gruyter & Co., Berlin, 2000.

[Ne08] —, *A complex semigroup approach to group algebras of infinite dimensional Lie groups*, Semigroup Forum **77** (2008), 5–35

[Ne13] —, *Unitary representations of unitary groups*, to appear in "Lie theory workshops", Eds. G. Mason, I. Penkov, J. Wolf, Progress in Math., Birkhäuser; arXiv:math.RT:1308.1500

[Nes13] Nessonov, N. I., *Schur–Weyl duality for the unitary groups of II_1-factors*, Preprint, arXiv.RT:1312.0824

[Ok66] Okayasu, T., *On the tensor products of C^*-algebras*, Tôhoku Math. J. (2) **18** (1966), 325–331

[Ol78] Olshanski, G. I., *Unitary representations of the infinite-dimensional classical groups* $\mathrm{U}(p,\infty)$*,* $\mathrm{SO}_0(p,\infty)$*,* $\mathrm{Sp}(p,\infty)$*, and of the corresponding motion groups*, Functional Anal. Appl. **12:3** (1978), 185–195

[Pau02] Paulsen, V., "Completely Bounded Maps and Operator Algebras", Cambridge Studies in Advanced Mathematics **78**, CUP, Cambridge, 2002.

[RSzN72] Riesz, F., and B. Sz.-Nagy, "Leçons d'analyse fonctionnelle", Sixième édition, Akadémiai Kiadó, Budapest, 1972.

[Seb86] Sebestyén, Z., *States and *-representations II*, Period. Math. Hungar. **17** (1986), no. 4, 247–254.

[St55] Stinespring, W. F., *Positive functions on C^*-algebras*, Proc. Amer. Math. Soc. **6** (1955), 211–216

[Tak02] Takesaki, M., "Theory of Operator Algebras. I", Encyclopaedia of Mathematical Sciences, 124. Operator Algebras and Non-commutative Geometry **5**, Springer-Verlag, Berlin, 2002

[Tr67] Treves, F., "Topological Vector Spaces, Distributions, and Kernels," Academic Press, New York, 1967.